

---

# Chapter 1

## Introduction

## Overview

This almanac contains information about current and historical air quality and emissions in California. In addition, forecasted emissions are presented. This document is a reference for anyone interested in air quality and emissions for criteria pollutants (NO<sub>x</sub>, ROG, CO, etc.) and toxic air contaminants (TACs). When using this information, please remember that the air quality and emission values are a snapshot of data at a particular point in time. This edition of the almanac is a year 2004 snapshot of the air quality and emission inventory databases. It is important to keep in mind that emission and air quality data can change over time. For example, emission data may be revised to reflect improved estimation methods, and air quality data may be changed because of corrections or additions of data.

The information in this document is based on data maintained in ARB's emission and air quality databases. The emission and human population estimates are presented at five-year intervals from 1975 to 2020. Data for vehicle miles traveled (VMT) are also provided at five-year increments, beginning with the year 1980. The air quality statistics in this almanac are for the period 1984 to 2003 for ozone, carbon monoxide (CO), nitrogen dioxide (NO<sub>2</sub>), sulfur dioxide (SO<sub>2</sub>), and lead. In addition, available 2004 statistics for ozone are also included for the five major air basins. PM<sub>10</sub> monitoring did not begin until 1988 and PM<sub>2.5</sub> monitoring until 1999. Therefore, PM<sub>10</sub> data cover the years 1988 to 2003, and PM<sub>2.5</sub> data cover the period 1999 through 2003. Air quality monitoring for TACs began in 1990; therefore, data for TACs covers the period 1990 through 2003.

## What's New in the 2005 Almanac?

- CD-ROM inside the back cover
- Web resources section (Chapter 1)
- Climate change discussion (Chapter 1)
- 8-Hour ozone statistics (Chapter 2, Chapter 4, Appendix A, and Appendix B)
- Statewide ammonia emissions table and discussion (Chapter 2)
- Statewide criteria pollutant emissions maps (Chapter 2)
- New federal designation maps (Chapter 2)
- PM<sub>2.5</sub> designation maps (Chapter 2)
- Emission, VMT/population, and diesel risk forecasts to 2020 (Chapter 3, Chapter 4, Appendix A, and Appendix D)
- Natural source emissions (Appendix E)

---

## Organization

This document is divided into five chapters and six appendices that include information, maps, graphs, and tabular data. Chapter 1 contains introductory material. Chapters 2 through 4 and Appendices A and B provide information on the most important criteria pollutants for which health-based ambient air quality standards have been established. Chapter 5 and Appendix C provide information on TACs. Appendix D includes information on population and vehicle miles traveled (VMT), and Appendix E contains information on natural emissions. In addition to this information, Appendix F provides lists of the figures and tables included in Chapters 1 through 5.

To help the reader navigate through the document, a short summary of each chapter and appendix is provided below:

- ◆ **Chapter 1** contains introductory material designed to help the reader better understand the remaining chapters. Included is information about data interpretation, emission estimation, air quality monitoring, the State and national standards, web resources, area designations for the State and national standards, and toxic air contaminants. A glossary of terms used in the Almanac as well as a list of air pollution contacts is provided at the end of this chapter.
- ◆ **Chapter 2** includes current emissions for NO<sub>x</sub>, ROG, PM<sub>10</sub>, PM<sub>2.5</sub>, CO, and NH<sub>3</sub> and air quality data for ozone, PM<sub>10</sub>, PM<sub>2.5</sub>, and CO for each air basin. The emission data also consists of lists of the State's highest emitting facilities. Information is included on how air quality in California compares to other parts of the country.
- ◆ **Chapter 3** provides historical emission and air quality trends from a statewide perspective. Statewide emission and air quality trends for ozone, PM<sub>10</sub>, PM<sub>2.5</sub>, CO, lead, and NO<sub>2</sub> are included. In addition, emission trends for oxides of sulfur (SO<sub>x</sub>) are included.
- ◆ **Chapter 4** provides historical emission and air quality trends for the State's five most populated regions. The pollutants covered are ozone, PM<sub>10</sub>, CO, and NO<sub>2</sub>.
- ◆ **Chapter 5** contains emission, air quality, and health risk information on TACs for the State as a whole and for five of California's most populated regions. The ten TACs, including diesel particulate matter (diesel PM), that pose the greatest risk in ambient (outdoor) air are covered. The air quality and health risk trends are based on ambient data collected during 1990 through 2003 (except for diesel PM, which is based on estimates of ambient concentrations).
- ◆ **Appendix A** includes more detailed emission data for NO<sub>x</sub>, ROG, PM<sub>10</sub>, PM<sub>2.5</sub>, and CO organized alphabetically, by air basin. Also included is a list of the highest emitting facilities in each air basin. Air quality data are provided for the criteria pollutants: ozone, PM<sub>10</sub>, PM<sub>2.5</sub>, CO, NO<sub>2</sub>, and SO<sub>2</sub>. Data are provided for all air basins and all counties (or county portions) within these air basins.
- ◆ **Appendix B** provides emission and air quality information similar to that found in Appendix A, but arranged by pollutant.

- ◆ **Appendix C** provides more detailed information on the ten TACs discussed in Chapter 5, including information on the emissions in each county and the air quality and health risk information for the individual sites where TAC concentrations are routinely measured.
- ◆ **Appendix D** provides tabulated information on surface area, population, and VMT for the State, each air basin, and for each county (or county portion) within the air basins.
- ◆ **Appendix E** provides emissions estimates for natural sources, including wildfires, vegetation (biogenic sources), and oil seeps (geogenic sources).
- ◆ **Appendix F** provides lists of the figures and tables included in Chapters 1 through 5.

*This almanac focuses on air emissions and air quality. The California Environmental Protection Agency (Cal/EPA) has developed a set of indicators to measure California's overall environmental health. The indicators cover all media, not just air, and help us understand the causes of environmental problems, the status of the environment, and the effectiveness of our environmental strategies. The data in this almanac are more detailed indicators of the State's air quality health, and in conjunction with Cal/EPA's indicators, provide a continuum of information from detailed air quality trends to California's overall environmental health. The most recent set of Cal/EPA indicators are available at <http://www.oehha.ca.gov/multimedia/epic/index.html>.*

---

## *California Facts and Figures*

California is truly a “Land of Contrasts.” The State offers a variety of physical features, including mountains, valleys, oceans, and deserts. In terms of size, California ranks third in the United States, after Alaska and Texas. California covers a total area of close to 160,000 square miles and is larger than many nations in the world today, including Great Britain, Japan, Italy, and Norway. Of California’s total area, about 152,000 square miles are land, and almost 8,000 square miles are water. The Pacific Ocean forms the western boundary of California, with a coastline more than 1,200 miles long. This is nearly equal to the combined Atlantic coastlines of Maine, New Hampshire, Massachusetts, Connecticut, Rhode Island, New York, and New Jersey.

California is blessed with a wide range of scenery and climates. The southern coastal areas enjoy a Mediterranean climate with the oak-studded hills and sunny beaches for which the State is famous. The northern coast is covered by fog-shrouded redwood forests. Inland lies the vast Central Valley with its millions of acres of cropland. The Sierra Nevada in the eastern half of California runs nearly two-thirds the length of the State. The Sierra includes the highest mountain in the continental United States, Mount Whitney, as well as the southernmost glacier in North America. Most of the southeastern portion of the State is desert, with sun-baked Death Valley, the lowest point in North America, lying only 60 miles from Mount Whitney. Further south are the scenic mountain ranges of the Mojave Desert.

To a large degree, California’s pleasant climate and abundance of relatively level land are the major features that have drawn people to the State. Since 1980, California’s population has increased about 54 percent, from nearly 24 million to nearly 37 million in the year 2004.

The increase in the average number of vehicle miles traveled (VMT) each day on our roadways has been even more dramatic. VMT has increased 120 percent, from about 389 million miles per day in 1980 to 857 million miles per day in 2004. With these dramatic increases in population and VMT have come tremendous challenges in controlling emissions to improve air quality.

## *Interpreting the Emission and Air Quality Statistics*

### *Interpreting Criteria Pollutant Emission and Air Quality Statistics.*

A number of pollutant trends are presented in this almanac. Emission and air quality trends for the same pollutant are usually highly correlated. In some cases, however, the two trends may differ, at least in terms of the rate of increase or decrease. The comparison of emission trends to air quality trends is complex, and a number of confounding factors can affect the resulting trends, such as the impacts of transported ozone and particulate matter from one area to another. An area can show a stable (or flat) emission trend because local emission growth offsets the reductions achieved through technology, but this same area may show an improvement in air quality because ambient concentrations reflect the impact of transport from a region that has improved. Other factors that can affect air quality are meteorology, which can cause large differences from year to year, and changes in monitoring sites (both site closures and the establishment of new sites). In addition, the emission data and some air quality statistics are based on estimates. These estimates use the best available methods, however, they embody some degree of uncertainty. All of these factors should be kept in mind when using and interpreting the trends.

Emission inventory trends make use of historical emission inventory data and projections based on expectations of future economic and population growth and emission controls. The historical emission inventory data in this almanac were updated to reflect improvements in emission inventory methodologies. The future year projections for stationary and areawide sources were developed using the California Emission Forecasting System (CEFS) model assuming a 2004 base year and California specific economic projections. These economic projections were prepared by E.H. Pechan and Associates and reflect information provided by local air districts. The stationary

source emission forecasts reflect control measure information received from local air districts as of September 2004. Future year emission projections for on and off road vehicles were developed using the ARB EMFAC 2002 version 2.2 and OFFROAD models, respectively. State Implementation Plan (SIP) and conformity inventory forecasts may differ from the forecasts presented in this almanac. For more information on these forecasts, please see the ARB SIP web page at [www.arb.ca.gov/sip/siprev1.htm](http://www.arb.ca.gov/sip/siprev1.htm).

In general, the criteria pollutant air quality trends in this almanac represent data that have been summarized from a network of monitoring sites to characterize the air quality in a particular region (for example, a county or air basin). Whenever data are summarized, the resulting statistics may be influenced by a number of factors, including the number of monitoring sites in operation and the completeness of the data. To help in interpreting the air quality trends, the ARB has included information on the time periods for which air quality data are available for different pollutants at sites in California and Baja, Mexico in its publication titled: “*California State and Local Air Monitoring Network Plan - 2003*” (October 2003). This report is available on the web at [www.arb.ca.gov/aqd/namlams/namlams.htm](http://www.arb.ca.gov/aqd/namlams/namlams.htm), or from the ARB’s Planning and Technical Support Division by calling (916) 322-5350.

A number of air quality statistics or indicators are used in this document. In general, 1-hour, 8-hour, and 24-hour concentrations reflect measured values and can be summarized by day, season or year. These data are also used to determine the number of days in which State or federal standards were exceeded. For the most part, this almanac provides data summarized as annual values. In contrast to measured

values, the peak indicators are calculated from the measured data. The peak indicator is used throughout the almanac for air quality trends for State standards. It represents the maximum concentration expected to be exceeded no more than once per year, on average based on the distribution of the data for each monitoring site. Because it is based on a robust statistical calculation using three years of data, it is relatively stable, thereby providing a trend indicator that is not highly influenced by year-to-year changes in weather. Finally, it is important to point out that the calculated number of days above the State and national PM<sub>10</sub> and PM<sub>2.5</sub> standards differ from other pollutants in that they are statistically derived from the measured data. This is because monitoring does not occur every day as with other criteria pollutants.

***Interpreting the Toxic Air Contaminant Emission and Air Quality Statistics.*** This almanac includes a number of statistics for the ten toxic air contaminants that generally pose the greatest ambient (outdoor) health risk to the public. These statistics are based on data collected by the ARB. TAC air quality data are also collected by the local air districts and for special studies. However, for consistency, only data collected by the ARB are included here. In general, TAC concentrations are sampled once every twelve days, for an average of two to three samples per month. Currently, the TAC sampling network comprises 18 sites located throughout California. The ARB originally established the network to measure the presence of TACs in the ambient air. The measured concentrations are generally used to represent average statewide concentrations and health risk. However, it is important to remember that concentrations can vary from one location to another. As a result, local concentrations and risks may be either higher or lower than the average values. Because the summary statistics are based on limited data, this information should be used with caution. The ARB has also been involved in efforts to better characterize local or community-wide exposures, and more information is available at [www.arb.ca.gov/ch/programs/sb25/sb25.htm](http://www.arb.ca.gov/ch/programs/sb25/sb25.htm).

A number of TAC emissions and air quality trends are presented in this almanac. Numerous factors influence the ambient measurements, and a number of assumptions are embodied in the summary statistics. The most important factors are summarized below. In contrast to the criteria pollutants, the control program for TACs is based on a risk assessment and risk management approach.

This almanac includes emissions data and ambient concentrations and information on the ten TACs that pose the greatest ambient risk in California: acetaldehyde, benzene, 1,3-butadiene, carbon tetrachloride, hexavalent chromium, formaldehyde, methylene chloride, *para*-dichlorobenzene, perchloroethylene, and diesel particulate matter (diesel PM). The determination of risk for the first nine TACs is based on ambient monitoring data while risk for diesel PM is based on estimates of ambient concentrations

It is important to note that the TAC information presented in this almanac reflects only those compounds for which data are available. There may be other compounds that pose a substantial risk, but have not been identified as a concern, or do not have data available. One example is dioxins, which may pose a substantial risk, but for which ambient air quality data are generally not available. Furthermore, the air quality and health risk information represents general population exposures. Therefore, the data may not provide information on localized impacts, often referred to as near-source exposures. The ARB has participated in several studies to address localized impacts and community health issues. Information from special studies in support of the community health program (SB25) and other studies are available at [www.arb.ca.gov/ch/programs/sb25/sb25.htm](http://www.arb.ca.gov/ch/programs/sb25/sb25.htm). The Web Resources section provides information on how to access this information.

The toxics emission inventory for 2004 represents the most currently available inventory compiled by the ARB staff. The toxic emissions for stationary sources include emissions data from the AB 2588 Air Toxics “Hot Spots” Program. For all source categories associated with

diesel fuel combustion, all particulate matter or “PM” emitted from these sources was considered “diesel PM.” The areawide source emissions were estimated by either the local air districts or the ARB staff. These toxic emission estimates were developed by speciating criteria emissions. Emission estimates for the other mobile source categories are primarily from ARB’s OFFROAD model, speciated for toxics. For the categories not currently included in the model, the emission estimates have been developed by either local air districts or ARB staff. Local air districts may also provide estimates for categories usually developed by ARB staff. Finally, the on-road mobile source emission estimates are based on the current model, EMFAC 2002, version 2.2. Again, the emission estimates have been speciated for toxics.

It is important to note that the health risk estimates reflect the estimated number of excess cancer cases per million people exposed over a 70-year period, and only for the ten compounds considered. In addition, the risk estimates are uncertain and actual health risk may be higher or lower than reported here. A number of factors add to the uncertainty, including the assumptions of the underlying risk factors, the assumption of a constant 70-year exposure, measurement biases and uncertainties, and the absence of ambient air quality data for diesel PM, dioxins, and other TACs that may pose a substantial health risk. However, the data are very useful for comparing relative health risks for the ten compounds considered (e.g., comparing the level of health risk for one compound or area relative to another).

While most of the TACs have some missing data caused by sampling or analysis problems, several TACs show substantial gaps in their data record. The limited sampling schedule only allows for two or three samples to be collected at each site during any month. In order to calculate a valid annual average (a mean of monthly means), each month during the year must have at least one valid measurement. Therefore, if there are no valid data in any given month, data for the

year will appear to be missing, even though some data may be available. Additionally, for TACs that have measurements below the limit of detection (LOD) of the instruments, we assign a value of one-half the LOD to these measurements when estimating an annual average.

In addition to missing data, there have been several site changes since the TAC network began operating. In several cases, the site changes occurred during the middle of a year. Because the site-by-site statistics presented in this almanac do not combine concentrations measured at different sites, an annual average for the year during which the site change occurred will be missing. Since all of the valid monthly means from each site are included in the air basin or statewide annual average, the site changes may lead to some variation in year-to-year statistics. In particular, the average health risk estimates may include a varying number of compounds and sites. Therefore, they may not be directly comparable from one year to the next. Site changes in each of the five major air basins are described in Chapter 5. A summary of the data record for all the monitoring sites and toxic air contaminants is available on ARB’s web site at [www.arb.ca.gov/aqd/toxics/toxics.html](http://www.arb.ca.gov/aqd/toxics/toxics.html). In addition, information about specific gaps in the TAC data is available from the ARB’s Monitoring and Laboratory Division at (916) 445-3742.

Finally, it is important to note that the concentrations and health risk estimates presented in this almanac are based on ambient outdoor measurements. They do not account for any indoor exposures to TACs. However, the indoor exposures can contribute significantly to individual health risk.

---

## *Meteorology's Role in Air Quality*

Air quality trends for a 20-year period are presented in this almanac. These trends reflect the progress that has been achieved as a result of our long history of emission control programs. Other factors such as meteorology (weather) and terrain can influence air quality trends. Meteorology can cause year-to-year changes in air quality in which the benefits of emission reductions can be masked. This is the reason why we focus on long-term trends in this almanac.

Ozone is formed in the atmosphere from emissions of oxides of nitrogen (NO<sub>x</sub>) and reactive organic gases (ROG) through a series of complex chemical reactions. Hot sunny days, with stagnant air and little dispersion (or mixing), are conditions that are most conducive to the formation of ozone. The highest ozone concentrations occur during the hot sunny days of the summer months. Also, winds can transport ozone or ozone precursors from one region to another, contributing to air quality problems downwind of the source regions.

As mentioned, California's terrain also plays a role in the formation of ozone. Mountains, such as the Sierra Nevada Mountains in the San Joaquin Valley and the San Bernardino Mountains in the Los Angeles area, can form a barrier that prevents ozone from dispersing.

Likewise, particulate matter (PM) can vary from year to year due to meteorology and terrain. Ambient PM is comprised of both directly emitted PM such as dust or soot, as well as PM formed in the atmosphere from the reactions of precursor gases, known as secondary PM. In some areas, windy days can result in high PM concentrations due to windblown dust. In other areas, cold temperatures, high humidity, and stagnant air can be conducive to the buildup of secondary PM. The amount and frequency of rainfall can also affect PM con-

centrations. For example, PM exceedances in the South Coast and San Joaquin Valley commonly occur in the winter months under these types of meteorological conditions.

Meteorological data can help determine how one year compares to the long-term average conditions (i.e., to the climate). Certain weather parameters are highly correlated to air quality, including temperature, the amount of sunlight, and the type of winds at the surface and above the surface. For example, higher horizontal wind speeds and greater vertical mixing (a function of how temperature changes above the surface) generally lead to lower air pollution levels.

Some years have fewer days with conditions that are conducive to air pollution. One of the important factors is improved mixing conditions. For ozone, improved mixing conditions result from a number of factors, including, more rain, cooler temperatures, or a change in the location of the normal summertime high-pressure area resulting in more days with more winds that allow for ozone to be dispersed. Greater dispersion can also result from a sea breeze which is more developed than in a 'normal' year. These subtle changes can result in less summertime ozone formation and/or in the dispersion or dilution of ozone or emissions that form ozone.

The Web Resources Section provides information on how to access sources of meteorological data. Sources such as ARB's real-time air quality and meteorological information system (AQMIS2) allow access to various wind parameters including wind speed/direction, temperature, humidity, and visibility.

## Sources of Emissions in California

California is a diverse state with many sources of air pollution. To estimate the sources and quantities of pollution, the ARB, in cooperation with local air districts and industry, maintains an inventory of California emission sources. Sources are subdivided into four major emission categories: stationary sources, area-wide sources, mobile sources, and natural sources.

Stationary source emissions are based on estimates made by facility operators and local air districts. Emissions from specific facilities can be identified by name and location. Area-wide emissions are estimated by ARB and local air district staffs. Emissions from area-wide sources may be either from small individual sources, such as residential fireplaces, or from widely distributed sources that cannot be tied to a single location, such as consumer products and dust from unpaved roads. Mobile source emissions are estimated by ARB staff with assistance from districts and other government agencies. Mobile sources include on-road cars, trucks, and buses and other sources such as boats, off-road recreational vehicles, aircraft, and trains. Natural sources are also estimated by the ARB staff and the air districts. These sources include biogenic hydrocarbons, geogenic hydrocarbons, natural wind-blown dust, and wildfires.

For the inventoried emission sources, the ARB compiles emission estimates for both the criteria pollutants and toxic air contaminants. Chapters 2 through 4 and Appendices A and B focus on five criteria pollutants: ozone, particulate matter, carbon monoxide, nitrogen dioxide, and sulfur dioxide. Emissions related to these criteria pollutants include reactive organic gases (ROG), oxides of nitrogen ( $\text{NO}_x$ ), carbon monoxide (CO), oxides of sulfur ( $\text{SO}_x$ ), ammonia ( $\text{NH}_3$ ), particulate matter with an aerodynamic diameter of 10 microns or

smaller ( $\text{PM}_{10}$ ), and particulate matter with an aerodynamic diameter of 2.5 microns or smaller ( $\text{PM}_{2.5}$ ).

While some pollutants, such as CO, are directly emitted, others are formed in the atmosphere from *precursor emissions*. Such is the case with ozone, which is formed in the atmosphere when hydrocarbon and  $\text{NO}_x$  precursor emissions react in the presence of sunlight. Particulate matter (PM), which includes  $\text{PM}_{10}$  and  $\text{PM}_{2.5}$ , is a complex pollutant that can either be directly emitted or formed in the atmosphere from precursor emissions. PM can form in the atmosphere from the reaction of gaseous precursors such as  $\text{NO}_x$ , ROG,  $\text{SO}_x$ , and ammonia. Examples of directly emitted PM include dust and soot.

*Hydrocarbon* is a general term used to describe compounds comprised of hydrogen and carbon atoms. Hydrocarbons are classified as to how photochemically reactive they are: relatively reactive or relatively non-reactive. Non-reactive hydrocarbons consist mostly of methane, which in turn consists of a single carbon atom and four hydrogen atoms. Emissions of *Total Organic Gases* and *Reactive Organic Gases* are two classes of hydrocarbons measured for California's emissions inventory. TOG includes all hydrocarbons, both reactive and non-reactive. In contrast, ROG includes only the reactive hydrocarbons.

In addition to the information about the criteria pollutants, Chapter 5 and Appendix C focus on the ten TACs that pose the greatest potential health risk, primarily based on statewide ambient air quality data. These ten TACs are: acetaldehyde, benzene, 1,3-butadiene, carbon tetrachloride, hexavalent chromium, *para*-dichlorobenzene, formaldehyde, methylene chloride, perchloroethylene, and diesel particulate matter. Excluding diesel particulate matter, the remaining nine TACs represent over 95 percent of the potential health risk as measured

---

through the statewide TAC air monitoring network. Although diesel particulate matter is not currently monitored, emissions and modeled ambient concentrations indicate that diesel particulate matter has a higher health risk than the other nine compounds combined.

## *Air Quality Monitoring*

Meteorology acts on the emissions released into the atmosphere to produce pollutant concentrations. These airborne pollutant concentrations are measured throughout California at air quality monitoring sites. The ARB operates a statewide network of monitors. Data from this network are supplemented with data collected by local air districts, other public agencies, and private contractors. As shown in Figure 1-1, there are more than 250 criteria pollutant monitoring sites in California. Currently, the ARB also monitors ambient concentrations of toxic air contaminants (TACs) at 18 of these sites. In addition to the California sites, a few monitoring sites are located in Mexico. These sites were established in cooperation with the United States Environmental Protection Agency (U.S. EPA) and the Mexican government to monitor the cross-border transport of pollutants and pollutant precursors. Each year, more than ten million air quality measurements from all of these sites are collected and stored in a comprehensive air quality database maintained by the ARB. To ensure the integrity of the data, the ARB routinely conducts audits and reviews of the monitoring instruments and the resulting data.




Figure 1-1

## California Air Basins

California contains a wide variety of climates, physical features, and emission sources. This variety makes the task of improving air quality complex, because what works in one area may not be effective in another area. To better manage common air quality problems, California is divided into 15 air basins, as shown in Figure 1-2 and Table 1-1. The ARB established the initial air basin boundaries during 1968.

An air basin generally has similar meteorological and geographical conditions throughout. To the extent possible, the air basin boundaries follow along political boundary lines and are defined to include both the source area and the receptor area. However, air masses can move freely from basin to basin. As a result, pollutants such as ozone and particulate matter can be transported across air basin boundaries, and interbasin transport is a reality that must be dealt with in air quality programs. Although established in 1968, the air basin boundaries have been changed several times over the years, to provide for better air quality management.



Figure 1-2

## List of Counties in Each Air Basin

### Great Basin Valleys Air Basin

- Alpine County
- Inyo County
- Mono County

### Lake County Air Basin

- Lake County

### Lake Tahoe Air Basin

- El Dorado County (portion)
- Placer County (portion)

### Mojave Desert Air Basin

- Kern County (portion)
- Los Angeles County (portion)
- Riverside County (portion)
- San Bernardino County (portion)

### Mountain Counties Air Basin

- Amador County
- Calaveras County
- El Dorado County (portion)
- Mariposa County
- Nevada County
- Placer County (portion)
- Plumas County
- Sierra County
- Tuolumne County

### North Central Coast Air Basin

- Monterey County
- San Benito County
- Santa Cruz County

### North Coast Air Basin

- Del Norte County
- Humboldt County
- Mendocino County
- Sonoma County (portion)
- Trinity County

### Northeast Plateau Air Basin

- Lassen County
- Modoc County
- Siskiyou County

### Sacramento Valley Air Basin

- Butte County
- Colusa County
- Glenn County
- Placer County (portion)
- Sacramento County
- Shasta County
- Solano County (portion)
- Sutter County
- Tehama County
- Yolo County
- Yuba County

---

## List of Counties in Each Air Basin

### Salton Sea Air Basin

- Imperial County
- Riverside County (portion)

### San Diego Air Basin

- San Diego County

### San Francisco Bay Area Air Basin

- Alameda County
- Contra Costa County
- Marin County
- Napa County
- San Francisco County
- San Mateo County
- Santa Clara County
- Solano County (portion)
- Sonoma County (portion)

### San Joaquin Valley Air Basin

- Fresno County
- Kern County (portion)
- Kings County
- Madera County
- Merced County
- San Joaquin County
- Stanislaus County
- Tulare County

### South Central Coast Air Basin

- San Luis Obispo County
- Santa Barbara County
- Ventura County

### South Coast Air Basin

- Los Angeles County (portion)
- Orange County
- Riverside County (portion)
- San Bernardino County (portion)

Table 1-1 (continued)

## *Criteria Air Pollutants*

### California and National Ambient Air Quality Standards

Very simply, an ambient air quality standard is the definition of “clean air.” More specifically, a standard establishes the concentration above which the pollutant is known to cause adverse health effects to sensitive groups within the population, such as children and the elderly. Both the California and federal governments have adopted health-based standards for the *criteria pollutants*, which include but are not limited to ozone, particulate matter (PM<sub>10</sub> and PM<sub>2.5</sub>), and carbon monoxide. For most pollutants the State standards are more stringent than the national standards. The differences in the standards are generally explained by the different health effects studies considered during the standard-setting process and the interpretation of the studies. In addition, the State standards incorporate a margin of safety to protect sensitive individuals (a short list of the State and national ambient air quality standards can be found on page 20, while a complete list can be found on the ARB website at [www.arb.ca.gov/aqs/aqs.htm](http://www.arb.ca.gov/aqs/aqs.htm)). In general, the air quality standards are expressed as a measure of the amount of pollutant per unit of air. For example, the particulate matter standards are expressed as micrograms of particulate matter per cubic meter of air ( $\mu\text{g}/\text{m}^3$ ).

## Ozone

Ozone, a colorless gas which is odorless at ambient levels, is the chief component of urban smog. Ozone is not directly emitted as a pollutant, but is formed in the atmosphere when hydrocarbon and NO<sub>x</sub> precursor emissions react in the presence of sunlight. Meteorology and terrain play major roles in ozone formation. Generally, low wind speeds or stagnant air coupled with warm temperatures and cloudless skies provide the optimum conditions for ozone formation. As a result, summer is generally the peak ozone season. Because of the reaction time involved, peak ozone concentrations often occur far downwind of the precursor emissions. Therefore, ozone is a regional pollutant that often impacts a large area.

Ozone impacts lung function by irritating and damaging the respiratory system. In addition, ozone causes damage to vegetation, buildings, rubber, and some plastics. Recognizing the impacts of day-long exposure, the U.S. EPA promulgated a new 8-hour standard for ozone in 1997.

On April 15, 2004 U.S. EPA designated areas of the country that exceed the 8-hour ozone standard as nonattainment. The designations became effective on June 15, 2004 and incorporate air quality data for the years 2001-2003. The designations can be accessed at U.S. EPA's website at [www.epa.gov/ozonedesignations](http://www.epa.gov/ozonedesignations). These designations will trigger new planning requirements for the 8-hour standard.

**State Ozone Standard:**

0.09 ppm for 1 hour,  
not to be exceeded.

**National Ozone Standards:**

0.12 ppm for 1 hour,  
not to be exceeded more  
than once per year *and*  
0.08 ppm for 8 hours,  
not to be exceeded,  
based on the fourth highest  
concentration averaged  
over three years.

Table 1-2

## Particulate Matter (PM<sub>10</sub> and PM<sub>2.5</sub>)

Exposure to particulate matter aggravates a number of respiratory illnesses and may even cause early death in people with existing heart and lung disease. Both long-term and short-term exposure can have adverse health impacts. All particles with a diameter of 10 microns or smaller (PM<sub>10</sub>) are harmful. For comparison, the diameter of a human hair is about 50 to 100 microns. PM<sub>10</sub> includes the subgroup of finer particles with an aerodynamic diameter of 2.5 microns or smaller (PM<sub>2.5</sub>). These finer particles pose an increased health risk because they can deposit deep in the lung and contain substances that are particularly harmful to human health.

PM<sub>10</sub> is a mixture of substances that includes elements such as carbon and metals; compounds such as nitrates, sulfates, and organic compounds; and complex mixtures such as diesel exhaust and soil. These substances may occur as solid particles or liquid droplets. Some particles are emitted directly into the atmosphere. Others, referred to as secondary particles, result from gases that are transformed into particles through physical and chemical processes in the atmosphere.

In 1982, the ARB adopted 24-hour average and annual average PM<sub>10</sub> standards. National ambient air quality standards for PM<sub>10</sub> have been in place since 1987. However, California's PM<sub>10</sub> standards are more health-protective.

In June 2002, the ARB adopted recommendations to lower the level of the PM<sub>10</sub> annual standard from 30  $\mu\text{g}/\text{m}^3$  to 20  $\mu\text{g}/\text{m}^3$  and established a new annual PM<sub>2.5</sub> standard of 12  $\mu\text{g}/\text{m}^3$ . The ARB plans to review short-term PM exposure studies in the future to determine if the current State 24-hour PM standards are adequately protective of public health. Additional information on the State PM standards is

available on the ARB's website at [www.arb.ca.gov/research/aaqs/std-rs/std-rs.htm](http://www.arb.ca.gov/research/aaqs/std-rs/std-rs.htm).

The U.S. EPA promulgated new national ambient air quality standards for PM<sub>2.5</sub> in 1997 to complement the national PM<sub>10</sub> standards. In early 2004, the ARB transmitted recommendations for area designations for the national PM<sub>2.5</sub> standards to U.S. EPA. U.S. EPA is scheduled to issue final designations by the end of 2004, with the designations becoming effective in early 2005.

### State PM<sub>10</sub> Standards:

50  $\mu\text{g}/\text{m}^3$  for 24 hours

not to be exceeded *and*

20  $\mu\text{g}/\text{m}^3$  annual arithmetic mean,  
not to be exceeded.

### State PM<sub>2.5</sub> Standard:

12  $\mu\text{g}/\text{m}^3$  annual arithmetic mean,  
not to be exceeded.

### National PM<sub>10</sub> Standards:

150  $\mu\text{g}/\text{m}^3$  for 24 hours, not to be exceeded,  
more than once per year *and*

50  $\mu\text{g}/\text{m}^3$  annual arithmetic mean  
averaged over 3 years, not to be exceeded.

### National PM<sub>2.5</sub> Standards:

65  $\mu\text{g}/\text{m}^3$  for 24 hours based on the  
98<sup>th</sup> percentile concentration averaged  
over three years, not to be exceeded *and*

15  $\mu\text{g}/\text{m}^3$  annual arithmetic mean  
averaged over 3 years, not to be exceeded.

Table 1-3

---

## Carbon Monoxide

Carbon monoxide is a colorless and odorless gas that is directly emitted as a by-product of combustion. The highest concentrations are generally associated with cold stagnant weather conditions that occur during winter. In contrast to ozone, which tends to be a regional pollutant, CO problems tend to be localized.

Carbon monoxide is harmful because it is readily absorbed through the lungs into the blood, where it binds with hemoglobin and reduces the ability of the blood to carry oxygen. As a result, insufficient oxygen reaches the heart, brain, and other tissues. The harm caused by CO can be critical for people with heart disease (angina), chronic lung disease, or anemia, as well as for unborn children. Even healthy people exposed to high levels of CO can experience headaches, fatigue, slow reflexes, and dizziness. Health damage caused by CO is of greater concern at high elevations where the air is less dense, aggravating the consequences of reduced oxygen supply. As a result, California has a more stringent CO standard for the Lake Tahoe Air Basin.

**State CO Standards:**

20 ppm for 1 hour *and*

9.0 ppm for 8 hours,

neither to be exceeded.

6 ppm for 8 hours

(Lake Tahoe Air Basin only),

not to be equaled or exceeded.

**National CO Standards:**

35 ppm for 1 hour *and*

9 ppm for 8 hours,

neither to be exceeded more

than once per year.

Table 1-4

## Air Quality Standards

Pollutant	Averaging Time	California Standards <sup>1</sup>	Federal Standards <sup>2</sup>	
		Concentration <sup>3</sup>	Primary <sup>3,4</sup>	Secondary <sup>3,5</sup>
Ozone (O <sub>3</sub> )	1 Hour	0.09 ppm (180 µg/m <sup>3</sup> )	0.12 ppm (235 µg/m <sup>3</sup> ) <sup>6</sup>	Same as Primary Standard
	8 Hour	—	0.08 ppm (157 µg/m <sup>3</sup> ) <sup>6</sup>	
Respirable Particulate Matter (PM <sub>10</sub> )	24 Hour	50 µg/m <sup>3</sup>	150 µg/m <sup>3</sup>	Same as Primary Standard
	Annual Arithmetic Mean	20 µg/m <sup>3</sup>	50 µg/m <sup>3</sup>	
Fine Particulate Matter (PM <sub>2.5</sub> )	24 Hour	No Separate State Standard	65 µg/m <sup>3</sup>	Same as Primary Standard
	Annual Arithmetic Mean	12 µg/m <sup>3</sup>	15 µg/m <sup>3</sup>	
Carbon Monoxide (CO)	8 Hour	9.0 ppm (10 mg/m <sup>3</sup> )	9 ppm (10 mg/m <sup>3</sup> )	None
	1 Hour	20 ppm (23 mg/m <sup>3</sup> )	35 ppm (40 mg/m <sup>3</sup> )	
	8 Hour (Lake Tahoe)	6 ppm (7 mg/m <sup>3</sup> )	—	
Nitrogen Dioxide (NO <sub>2</sub> )	Annual Arithmetic Mean	—	0.053 ppm (100 µg/m <sup>3</sup> )	Same as Primary Standard
	1 Hour	0.25 ppm (470 µg/m <sup>3</sup> )	—	
Sulfur Dioxide (SO <sub>2</sub> )	Annual Arithmetic Mean	—	0.030 ppm (80 µg/m <sup>3</sup> )	—
	24 Hour	0.04 ppm (105 µg/m <sup>3</sup> )	0.14 ppm (365 µg/m <sup>3</sup> )	—
	3 Hour	—	—	0.5 ppm (1300 µg/m <sup>3</sup> )
	1 Hour	0.25 ppm (655 µg/m <sup>3</sup> )	—	—
Lead <sup>7</sup>	30 Day Average	1.5 µg/m <sup>3</sup>	—	—
	Calendar Quarter	—	1.5 µg/m <sup>3</sup>	Same as Primary Standard

- California standards for ozone, carbon monoxide (except Lake Tahoe), sulfur dioxide (1 and 24 hour), nitrogen dioxide, suspended particulate matter—PM<sub>10</sub>, PM<sub>2.5</sub>, and visibility reducing particles, are values that are not to be exceeded. All others are not to be equaled or exceeded. California ambient air quality standards are listed in the Table of Standards in Section 70200 of Title 17 of the California Code of Regulations.
- National standards (other than ozone, particulate matter, and those based on annual averages or annual arithmetic mean) are not to be exceeded more than once a year. The ozone standard is attained when the fourth highest eight hour concentration in a year, averaged over three years, is equal to or less than the standard. For PM<sub>10</sub>, the 24 hour standard is attained when the expected number of days per calendar year with a 24-hour average concentration above 150 µg/m<sup>3</sup> is equal to or less than one. For PM<sub>2.5</sub>, the 24 hour standard is attained when 98 percent of the daily concentrations, averaged over three years, are equal to or less than the standard. Contact U.S. EPA for further clarification and current federal policies.
- Concentration expressed first in units in which it was promulgated. Equivalent units given in parentheses are based upon a reference temperature of 25°C and a reference pressure of 760 torr. Most measurements of air quality are to be corrected to a reference temperature of 25°C and a reference pressure of 760 torr, ppm in this table refers to ppm by volume, or micromoles of pollutant per mole of gas.
- National Primary Standards: The levels of air quality necessary, with an adequate margin of safety to protect the public health.
- National Secondary Standards: The levels of air quality necessary to protect the public welfare from any known or anticipated adverse effects of a pollutant.
- New federal 8-hour ozone and fine particulate matter standards were promulgated by U.S. EPA on July 18, 1997. Contact U.S. EPA for further clarification and current federal policies.
- The ARB has identified lead and vinyl chloride as 'toxic air contaminants' with no threshold level of exposure for adverse health effects determined. These actions allow for the implementation of control measures at levels below the ambient concentrations specified for these pollutants.

Table 1-5

---

*(This page intentionally left blank)*

## California and National Area Designations

Both the California and federal governments use monitoring data to designate areas according to their attainment status for most of the pollutants with ambient air quality standards. The purpose of the designations is to identify those areas with air quality problems and thereby initiate planning efforts to make the air more healthful. There are three basic designation categories: nonattainment, attainment, and unclassified. In addition, the California (State) designations include a subcategory of the nonattainment designation, called nonattainment-transitional. The nonattainment-transitional designation is given to nonattainment areas that are making progress and nearing attainment.

A *nonattainment designation* indicates that the air quality violates an ambient air quality standard. Although a number of areas may be designated as nonattainment for a particular pollutant, the severity of the problem can vary greatly. For example, in two ozone nonattainment areas, the first area has a measured maximum concentration of 0.13 parts per million (ppm), while the second area has a measured maximum concentration of 0.23 ppm. While both areas are designated as nonattainment, it is obvious that the second area has a more severe ozone problem and will need a more stringent emission control strategy. To identify the severity of the problem and the extent of planning required, nonattainment areas are assigned a classification that is commensurate with the severity of their air quality problem (e.g., moderate, serious, severe).

In contrast to nonattainment, an *attainment designation* indicates that the air quality does not violate the established standard. Under the federal Clean Air Act, nonattainment areas that are redesignated as attainment must develop and implement maintenance plans designed to assure continued compliance with the standard.

Finally, an *unclassified designation* indicates that there are insufficient data for determining attainment or nonattainment. The U.S. EPA combines unclassified and attainment into one designation for ozone and carbon monoxide. More detailed information on the area designation categories can be found on the ARB's website at [www.arb.ca.gov/desig/desig.htm](http://www.arb.ca.gov/desig/desig.htm).

## Ozone - State Area Designations

Some rural and northern coastal areas of California are designated as attainment for the State ozone standard. Last year, two new areas attained the State standard: San Luis Obispo County and the Northern Sonoma Air Quality Management District. However, most of the rest of the State, including all of the major urban areas, have ozone concentrations that violate the State standard, and therefore, are designated as nonattainment. Although few areas have made sufficient progress to be redesignated as attainment for the State ozone standard, ozone precursor emissions continue to decline throughout California. As a result, air quality is improving, and more areas should eventually qualify for attainment status, as precursor emissions continue to be reduced statewide.



Figure 1-3

## Ozone - National 1-Hour Area Designations

Similar to the State designations, most of the major urban areas in California are designated as nonattainment for the national 1-hour ozone standard. In contrast to the State designations, there are only two designation categories for the national ozone standards: attainment/unclassified and nonattainment.

Ozone air quality in Butte, northern Sutter, Yuba, and Ventura counties, and the San Francisco Bay Area Air Quality Management District now meets the national 1-hour ozone standard, but these areas have not yet been officially redesignated as attainment. Eastern Kern county has been officially redesignated as attainment/unclassified. All the remaining areas of California are designated as attainment/unclassified. Ambient ozone concentrations in these areas do not violate the national 1-hour standard. However, a number of these areas violate the new 8-hour national ozone standard.

U.S. EPA has indicated that it intends to revoke the 1-hour standard by June 15, 2005. However, nonattainment areas are still required to complete their 1-hour planning requirements or submit early 8-hour ozone plans. For more information refer to the U.S. EPA website at [www.epa.gov/ttn/naaqs/ozone/o3imp8hr/finalrule.html](http://www.epa.gov/ttn/naaqs/ozone/o3imp8hr/finalrule.html).



Figure 1-4

## Ozone - National 8-Hour Area Designations

On April 15, 2005, the U.S. EPA made first time designations for the national 8-hour ozone standard. There are 15 nonattainment areas in California, including all of the five largest urban areas in California. In addition, a number of smaller counties and rural areas exceed the standard. An area violates the federal 8-hour ozone standard if the calculated fourth highest 8-hour concentration averaged over a three-year period exceeds the attainment level of the standard at any monitoring site in the region.

The designations became effective on June 15, 2004, and triggered new planning requirements. State Implementation Plans will be due to the U.S. EPA by 2007. Until then, all of our programs designed to achieve the national 1-hour standard will bring us closer to attaining the national 8-hour standard. The U.S. EPA will publish an implementation rule later this year that will identify many of the specific planning and control requirements for the 8-hour standard.




Figure 1-5

## PM<sub>10</sub> - State Area Designations

The majority of California is designated as nonattainment for the State PM<sub>10</sub> standards. Three counties in the northern half of the State remain unclassified, and one area, the Lake County Air Basin has been designated as attainment. In addition, Siskiyou County has been proposed to be redesignated as attainment.

PM<sub>10</sub> remains a widespread problem, and its causes are very diverse. Because of the variety of sources and the size and chemical make-up of the particles, the PM<sub>10</sub> problem can vary considerably from one area to the next. In addition, high PM<sub>10</sub> concentrations are seasonal, and the high season varies from area to area. For example, in some areas, windblown dust may contribute to high PM<sub>10</sub> concentrations in the summer and fall, while in other areas, high concentrations due to secondary particles may occur during the winter. As a result, two areas with similar PM<sub>10</sub> concentrations may have very different PM<sub>10</sub> problems, and multiple control strategies are needed to effectively deal with these problems.



Figure 1-6

## PM<sub>10</sub> - National Area Designations

In contrast to the State PM<sub>10</sub> designations, there are only two designation categories for the national PM<sub>10</sub> standards: attainment/unclassified and nonattainment. Areas designated as nonattainment for the national PM<sub>10</sub> standards are required to develop and implement plans designed to meet the standards. Although they are still designated as nonattainment, Sacramento, Mammoth Lakes, Trona (northwestern San Bernardino County), and the portion of San Bernardino County outside of the South Coast air basin now meet the national PM<sub>10</sub> standards.



Figure 1-7

## PM<sub>2.5</sub> - State Area Designations

Since California adopted the new PM<sub>2.5</sub> standard in 2002, this is the second year of area designations for the State PM<sub>2.5</sub> standard. Approximately half of California is designated as nonattainment for the State PM<sub>2.5</sub> standard, with Lake County Air Basin as the only attainment area. Additionally, Lake Tahoe and North Central Coast Air Basins are proposed to be designated attainment this year. Nonattainment areas include all of the major urban areas as well as a few rural areas. Secondary formation of PM<sub>2.5</sub> and particles directly emitted from combustion processes are major contributors to high PM<sub>2.5</sub> concentrations in these areas.

California's programs to reduce ozone, PM<sub>10</sub>, and diesel PM are helping reduce PM<sub>2.5</sub>. In addition, as required by new legislation enacted in 2003 (Senate Bill 656), ARB has assembled a list of measures that can be used by air districts to further reduce PM and PM precursor emissions, it is available on the web at [www.arb.ca.gov/pm/pmmeasures/pmmeasures.htm](http://www.arb.ca.gov/pm/pmmeasures/pmmeasures.htm).




Figure 1-8

## PM<sub>2.5</sub> - National Area Designations

The U. S. EPA is expected to promulgate first time area designations for PM<sub>2.5</sub> at the end of 2004. The San Joaquin Valley, South Coast, and San Diego air basins are expected to be designated nonattainment. These air basins include major urban areas as well as some rural areas. Reactions in the atmosphere from precursor gases emitted from combustion sources and direct particulate emissions from mobile sources and burning activities lead to high PM<sub>2.5</sub> concentrations in these areas. The remaining areas of the State are designated attainment/unclassified.

State Implementation Plans for PM<sub>2.5</sub> nonattainment areas are scheduled for submittal in early 2008. Until then, actions taken to reduce ozone, PM<sub>10</sub> and diesel PM will also help in reducing PM<sub>2.5</sub>.



Designations Expected December 2004

Figure 1-9

## Carbon Monoxide - State Area Designations

Currently, there is only one area in the state that does not meet the State CO standards: the City of Calexico, in Imperial County. California has made tremendous progress in reducing CO concentrations in the last ten years, during which a number of areas have been redesignated as attainment. Most recently, Los Angeles County qualified for attainment and will be considered for redesignation as attainment in early 2005. Much of the progress in reducing ambient CO is attributable to motor vehicle controls and the introduction of cleaner fuels.

With respect to the nonattainment area, the problem in Calexico is unique in that this area is probably impacted by emissions from Mexico. Additional studies are needed to determine the most effective control strategy for the Calexico area.



Figure 1-10

## Carbon Monoxide - National Area Designations

The U.S. EPA uses only two designation categories for CO: attainment/unclassified and nonattainment. All areas of California except the South Coast Air Basin are currently designated as attainment/unclassified for the national CO standards. However, the South Coast Air Basin now meets the requirements for attainment, although they have not been officially redesignated. Most CO is directly emitted by cars and trucks, and the ARB's motor vehicle controls should be sufficient to overcome the problem in the coming years. In addition, the City of Calexico, in Imperial County, has carbon monoxide concentrations that violate the national standards. There is a high likelihood that cross-border traffic contributes to the local CO problem in this area, and more study is needed to determine the most effective control strategy. The U.S. EPA has not acted to change this area's designation from attainment/unclassified to nonattainment.



Figure 1-11

## *Toxic Air Contaminants*

A toxic air contaminant or TAC is defined as an air pollutant which may cause or contribute to an increase in mortality or in serious illness, or which may pose a hazard to human health. TACs are usually present in minute quantities in the ambient air. However, their high toxicity or health risk may pose a threat to public health even at very low concentrations. In general, for those TACs that may cause cancer, there is no concentration that does not present some risk. In other words, there is no threshold level below which adverse health impacts may not be expected to occur. This contrasts with the criteria pollutants for which acceptable levels of exposure can be determined and for which the State and federal governments have set ambient air quality standards.

The Air Resources Board's TAC program traces its beginning to the criteria pollutant program in the 1960s. For many years, the criteria pollutant control program has been effective at reducing TACs since many volatile organic compounds and PM constituents are also TACs. During the 1980s, the public's concern over toxic chemicals heightened. As a result, citizens demanded protection and control over the release of toxic chemicals into the air. In response to public concerns, the California legislature enacted a 1983 law governing the release of TACs. This law charges the Air Resources Board with the responsibility for identifying substances as TACs, setting priorities for control, adopting control strategies, and promoting alternative processes. To date, the ARB has designated nearly 192 compounds as TACs. Additionally, the ARB has implemented control strategies for a number of compounds that pose high risk and show potential for effective control.

The majority of the estimated health risk from TACs can be attributed to a relatively few compounds, the most important being PM

from diesel-fueled engines (diesel particulate matter or diesel PM). Diesel PM differs from other TACs in that it is not a single substance, but rather a complex mixture of hundreds of substances. Although diesel PM is emitted by diesel-fueled internal combustion engines, the composition of the emissions will vary depending on engine type, operating conditions, fuel composition, lubricating oil, and whether an emission control system is present. Unlike the other toxic air contaminants, no ambient monitoring data are available for diesel PM because no routine measurement method currently exists. However, the ARB has made preliminary concentration estimates for the State and its 15 air basins using a PM-based exposure method. The method uses the ARB emission inventory's PM<sub>10</sub> database, ambient PM<sub>10</sub> monitoring data, and the results from several studies on chemical speciation of ambient data. These data were used, along with receptor modeling techniques, to estimate outdoor concentrations of diesel PM. Details on the method and the resulting estimates for individual air basins can be found in the ARB report entitled: "*Proposed Identification of Diesel Exhaust as a Toxic Air Contaminant -- Appendix III Part A Exposure Assessment*," (April 1998). In addition to diesel PM, benzene and 1,3-butadiene are also significant contributors to overall ambient public health risk in California.

This almanac (Chapter 5 and Appendix C) includes information for ten TACs: acetaldehyde, benzene, 1,3-butadiene, carbon tetrachloride, hexavalent chromium, *para*-dichlorobenzene, formaldehyde, methylene chloride, perchloroethylene, and diesel PM. These ten compounds pose the greatest ambient risk based on air quality data, or concentration estimates in the case of diesel PM. Note that other TACs, for example dioxins, may also pose a significant health risk. However, sufficient air quality data are not yet available for these

compounds. In order to provide information on ambient levels of dioxins and dioxin-like compounds, the Air Resources Board is conducting the California Ambient Dioxin Air Monitoring Program (CADAMP). Ten sampling sites have been deployed in CADAMP, five in the San Francisco Bay Area, four in the South Coast Air Basin, and one in Sacramento. Monitoring at the ten sites will continue until December 2004 with over 400 samples to be analyzed. More information on the dioxin monitoring program, including results, can be found at [www.arb.ca.gov/aaqm/qmosopas/dioxins/overview.htm](http://www.arb.ca.gov/aaqm/qmosopas/dioxins/overview.htm).

Most of the TAC data in this almanac were obtained from monitors operated by the ARB. The majority of the information is presented on a pollutant-by-pollutant basis, with a focus on cancer risk. The available data represent average population exposures and may not represent the health risk near local sources. Localized impacts may involve exposure to different toxic air contaminants or to higher or lower concentrations than those represented by the ambient monitoring data. ARB participated in several studies to address localized impacts and community health issues to learn which communities are the most impacted and who in those communities are the most vulnerable. More information on these studies is available on the web at [www.arb.ca.gov/ch/ch.htm](http://www.arb.ca.gov/ch/ch.htm).

The ARB has substantially increased its knowledge about TACs in the last 15 years, and the data indicate that control efforts have been effective in reducing public exposures and associated health risks. The future gradual phase-in of control strategies will likely continue to result in lower exposures for California's citizens. In the interim, work continues on identifying toxic substances and developing a better understanding of the risks they pose. Health experts still have only a limited knowledge of the mechanisms by which many toxic substances harm the body, and there is still much work to be done in researching health effects and quantifying cancer risks. Cooperative strategies between the ARB, businesses, and other State, local, and

federal agencies will be a major focus of future control efforts.

Additional information on TACs may be found on the ARB website at [www.arb.ca.gov/toxics/toxics.htm](http://www.arb.ca.gov/toxics/toxics.htm). Detailed information on the health effects of these pollutants, as well as many other toxic air contaminants, can be found in a report entitled: "*Toxic Air Contaminant Identification List-Summaries*." This report, dated September 1997, is available from the ARB Public Information Office and on the web at [www.arb.ca.gov/toxics/tac/intro.htm](http://www.arb.ca.gov/toxics/tac/intro.htm).



Figure 1-12

## Climate Change

The earth's climate has been warming for the past century. It is believed that this warming trend is related to the release of certain gases into the atmosphere. The greenhouse gases (GHG) include carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>), nitrous oxide (N<sub>2</sub>O), and hydrofluorocarbons. Climate research has identified other greenhouse agents that can drive climate change, particularly tropospheric ozone, and atmospheric aerosols (particles containing sulfate, black carbon or other carbonaceous compounds).

Greenhouse gases absorb infrared energy that would otherwise escape from the Earth. As the infrared energy is absorbed, the air surrounding the earth is heated. An overall warming trend has been recorded since the late 19th century, with the most rapid warming occurring over the past two decades. The 10 warmest years of the last century all occurred within the last 15 years. It appears that the decade of the 1990s was the warmest in human history.

It is a fact that human activities have increased the atmospheric abundance of greenhouse gases. There are uncertainties as to exactly what the climate changes will be in various local areas of the Earth, and what the effects of clouds will be in determining the rate at which the mean temperature will increase. There are also uncertainties associated with the magnitude and timing of other consequences of a warmer planet: sea level rise, spread of certain diseases out of their usual geographic range, the effect on agricultural production, water supply, sustainability of ecosystems, increased strength and frequency of storms, extreme heat events, air pollution episodes, and the consequence of these effects on the economy. Already, some of these effects have been seen in California.

The United States has the highest emissions of greenhouse gases of any nation on Earth. California's transportation sector is the single largest contributor of GHGs in the State, producing close to 60% of all such emissions. In the absence of controls, the State's inventory of greenhouse gases would mirror the growth in population; transportation and land-use trends in California would continue to increase greenhouse gas production.

California has been studying the impacts of climate change since 1988, when AB4420 was approved. This legislation directed the California Energy Commission, in consultation with the Air Resources Board and other agencies, to study the implications of global warming on California's environment, economy, and water supply. The Energy Commission was also directed to prepare and maintain the state's inventory of GHG emissions. More information on the Energy Commission's climate change activities can be found on the web at [www.energy.ca.gov/global\\_climate\\_change/index.html](http://www.energy.ca.gov/global_climate_change/index.html).

In 2002, recognizing that global warming would impact California, AB1493 was approved. That bill directed the ARB to adopt regulations to achieve the maximum feasible and cost-effective reduction of greenhouse gas emissions from motor vehicles. ARB staff's proposal implementing these regulations was approved by the Air Resources Board in September, 2004. These regulations will be reviewed and may be modified by the California Legislature. The average reduction of greenhouse gases from new California cars and light trucks will be about 22 percent in 2012 and about 30 percent in 2016, compared to today's vehicles. More information on ARB's Climate Change regulations can be found on the web at [www.arb.ca.gov/cc/cc.htm](http://www.arb.ca.gov/cc/cc.htm).

---

## *California Air Quality Regulation*

The responsibility for controlling air pollution in California is shared between 35 local air districts, the Air Resources Board, and the United States Environmental Protection Agency. The basic responsibilities of each of these entities are outlined below.

### **District Responsibilities:**

- Control and permit industrial pollution sources (such as power plants, refineries, and manufacturing operations) and widespread area-wide sources (such as bakeries, dry cleaners, service stations, and commercial paint applicators).
- Adopt local air quality plans and rules.

### **Air Resources Board Responsibilities:**

- Establish State ambient air quality standards.
- Adopt and enforce emission standards for mobile sources (except where federal law preempts ARB's authority), fuels, consumer products, and toxic air contaminants.
- Provide technical support to the local districts.
- Oversee local district compliance with State and federal law.
- Approve local air quality plans and submit State Implementation Plans to U.S. EPA.

### **United States Environmental Protection Agency Responsibilities:**

- Establish national ambient air quality standards.
- Set emission standards for mobile sources, including those sources under exclusive federal jurisdiction (like interstate trucks, aircraft, marine vessels, locomotives, and farm/construction equipment).

- Oversee State air programs as they relate to the Federal Clean Air Act.
- Approve State Implementation Plans.

## *List of Air Pollution Contacts*

### **Amador County Air Pollution Control District**

All of Amador County

(209) 257-0112

[www.amadorapcd.org](http://www.amadorapcd.org)

### **Antelope Valley Air Pollution Control District**

Northeast portion of Los Angeles County

(661) 723-8070

[www.avaqmd.ca.gov](http://www.avaqmd.ca.gov)

### **Bay Area Air Quality Management District**

All of Alameda, Contra Costa, Marin, Napa, San Francisco, San Mateo, and Santa Clara counties, western portion of Solano County, and southern portion of Sonoma County

(415) 749-5000

[www.baaqmd.gov](http://www.baaqmd.gov)

### **Butte County Air Quality Management District**

All of Butte County

(530) 891-2882

[www.bcaqmd.org](http://www.bcaqmd.org)

### **Calaveras County Air Pollution Control District**

All of Calaveras County

(209) 754-6504

[www.co.calaveras.ca.us/departments/env.html](http://www.co.calaveras.ca.us/departments/env.html)

### **Colusa County Air Pollution Control District**

All of Colusa County

(530) 458-0590

[www.colusanet.com/apcd](http://www.colusanet.com/apcd)

### **El Dorado County Air Pollution Control District**

All of El Dorado County

(530) 621-6662

[www.co.el-dorado.ca.us/emd/apcd](http://www.co.el-dorado.ca.us/emd/apcd)

### **Feather River Air Quality Management District**

All of Sutter and Yuba counties

(530) 634-7659

[www.fraqmd.org](http://www.fraqmd.org)

### **Glenn County Air Pollution Control District**

All of Glenn County

(530) 934-6500

[www.countyofglenn.net/Air\\_Pollution\\_Control/home\\_page.asp](http://www.countyofglenn.net/Air_Pollution_Control/home_page.asp)

### **Great Basin Unified Air Pollution Control District**

All of Alpine, Inyo, and Mono counties

(760) 872-8211

[gb1@greatbasinapcd.org](mailto:gb1@greatbasinapcd.org)

### **Imperial County Air Pollution Control District**

All of Imperial County

(760) 482-4606

[www.imperialcounty.net](http://www.imperialcounty.net)

### **Kern County Air Pollution Control District**

Eastern portion of Kern County

(661) 862-5250

[www.kernair.org](http://www.kernair.org)

---

**Lake County Air Quality Management District**

All of Lake County  
(707) 263-7000  
[www.lcaqmd.net](http://www.lcaqmd.net)

**Lassen County Air Pollution Control District**

All of Lassen County  
(530) 251-8110  
[lassenag@psln.com](mailto:lassenag@psln.com)

**Mariposa County Air Pollution Control District**

All of Mariposa County  
(209) 966-2220  
[www.mariposacounty.org/healthdepartment/index.html](http://www.mariposacounty.org/healthdepartment/index.html)

**Mendocino County Air Quality Management District**

All of Mendocino County  
(707) 463-4354  
[www.co.mendocino.ca.us/aqmd](http://www.co.mendocino.ca.us/aqmd)

**Modoc County Air Pollution Control District**

All of Modoc County  
(530) 233-6419  
[modapcd@hdo.net](mailto:modapcd@hdo.net)

**Mojave Desert Air Quality Management District**

Northern portion of San Bernardino County and eastern portion of Riverside County  
(760) 245-1661  
[www.mdaqmd.ca.gov](http://www.mdaqmd.ca.gov)

**Monterey Bay Unified Air Pollution Control District**

All of Monterey, San Benito and Santa Cruz counties  
(831) 647-9411  
[www.mbuapcd.org](http://www.mbuapcd.org)

**North Coast Unified Air Quality Management District**

All of Del Norte, Humboldt, and Trinity counties  
(707) 443-1580  
[www.ncuaqmd.org](http://www.ncuaqmd.org)

**Northern Sierra Air Quality Management District**

All of Nevada, Plumas, and Sierra counties  
(530) 274-9360  
[www.nccn.net/~nsaqmd](http://www.nccn.net/~nsaqmd)

**No. Sonoma County Air Pollution Control District**

Northern portion of Sonoma County  
(707) 433-5911  
[nsc@sonic.net](mailto:nsc@sonic.net)

**Placer County Air Pollution Control District**

All of Placer County  
(530) 889-7130  
[www.placer.ca.gov/airpollution/airpolut.htm](http://www.placer.ca.gov/airpollution/airpolut.htm)

**Sacramento Metro Air Quality Management District**

All of Sacramento County  
(916) 874-4800  
[www.airquality.org](http://www.airquality.org) or [www.sparetheair.com](http://www.sparetheair.com)

**San Diego County Air Pollution Control District**

All of San Diego County  
(858) 650-4650  
[www.sdapcd.co.san-diego.ca.us](http://www.sdapcd.co.san-diego.ca.us)

**San Joaquin Valley Unified Air Pollution Control District**

All of Fresno, Kings, Madera, Merced, San Joaquin, Stanislaus, and Tulare counties and western portion of Kern County  
(559) 230-6000  
[www.valleyair.org](http://www.valleyair.org)

**San Luis Obispo County Air Pollution Control District**

All of San Luis Obispo County

(805) 781-4AIR

[www.slocleanair.org](http://www.slocleanair.org)

**Santa Barbara County Air Pollution Control District**

All of Santa Barbara County

(805) 961-8800

[www.sbcapcd.org](http://www.sbcapcd.org)

**Shasta County Air Quality Management District**

All of Shasta County

(530) 225-5674

[www.co.shasta.ca.us/Departments/Resourcemgmt/drm/aqmain.htm](http://www.co.shasta.ca.us/Departments/Resourcemgmt/drm/aqmain.htm)

**Siskiyou County Air Pollution Control District**

All of Siskiyou County

(530) 841-4029

[ebeck@co.siskiyou.ca.us](mailto:ebeck@co.siskiyou.ca.us)

**South Coast Air Quality Management District**

Los Angeles County except for portion covered by Antelope Valley APCD, all of Orange County, western portion of San Bernardino County, and western portion of Riverside County

(909) 396-2000

[www.aqmd.gov](http://www.aqmd.gov)

**Tehama County Air Pollution Control District**

All of Tehama County

(530) 527-3717

[www.tehcoapcd.net](http://www.tehcoapcd.net)

**Tuolumne County Air Pollution Control District**

All of Tuolumne County

(209) 533-5693

[bsandman@co.tuolumne.ca.us](mailto:bsandman@co.tuolumne.ca.us)

**Ventura County Air Pollution Control District**

All of Ventura County

(805) 645-1400

[www.vcapcd.org](http://www.vcapcd.org)

**Yolo-Solano Air Quality Management District**

All of Yolo County and eastern portion of Solano County

(530) 757-3650

[www.ysaqmd.org](http://www.ysaqmd.org)

---

## *Milestones in California's Emission Control Programs*

### **Historical Milestones:**

- 1963:** First vehicle emission control in the country – positive crankcase ventilation required to reduce evaporative emissions.
- 1966:** First tailpipe emission standards for hydrocarbons (HC) and carbon monoxide (CO).
- 1971:** First oxides of nitrogen (NO<sub>x</sub>) standards for cars and light trucks.
- 1973:** First heavy-duty diesel truck standards.
- 1975:** Two-way catalytic converters first used to control HC and emissions from cars.
- 1976:** “Unleaded” gasoline first offered for sale, with reduced lead levels.  
Three-way catalyst first used to control NO<sub>x</sub>, HC, and CO emissions from cars.
- 1984:** California Smog Check program implemented to identify and repair ineffective emission control systems on cars and light-trucks.
- 1988:** California Clean Air Act is enacted, setting forth the framework for meeting State ambient air quality standards.
- 1992:** California’s reformulated gasoline introduced – reducing evaporative emissions, phasing out lead in gasoline, and requiring wintertime oxygenates to reduce CO formation.
- First consumer product regulations take effect, regulating HC emissions from aerosol antiperspirants and deodorants.
- 1993:** Cleaner diesel fuel launched, reducing emissions of diesel particulate matter, sulfur dioxide, and NO<sub>x</sub>.  
Regulations to limit HC emissions from consumer products such as hairspray, windshield washer fluid, and air fresheners take effect.
- 1994:** Low emission vehicle regulations to further reduce emissions from cars and light trucks take effect.
- 1996:** Cleaner burning gasoline debuts with emission benefits equivalent to removing 3.5 million cars from California roads.  
Regulations reducing HC emissions from spray paint take effect.
- 1998:** Tighter standards for California diesel trucks and buses take effect.  
Revamped Smog Check II program implemented.
- 1999:** ARB acted to phaseout MTBE in gasoline.
- 2000:** Tighter emission standards for off-road diesel equipment, such as tractors and generators, take effect nationwide.  
More stringent California standards for the small engines used in lawn and garden equipment take effect.  
Diesel Risk Reduction Plan adopted.
- 2001:** First California standards for large spark ignition off-road engines such as forklifts and pumps take effect.

More stringent standards for pleasure boats and personal watercraft sold in California begin.

Limits on HC emissions from products such as carpet and upholstery cleaners take effect.

**2002:** Emission standards for new heavy-duty diesel trucks are cut in half, nationwide.

**2003:** New emission standards for inboard marine engines sold in California take effect.

#### **Upcoming Milestones:**

**2004:** Regulations to further reduce emissions from cars (and require light-trucks and sport-utility vehicles to meet the same emission standards as cars) take effect in California.

MTBE in California gasoline is fully phased out.

Tighter standards for on-road motorcycles begin.

**2005:** Limits on HC emissions from paint removers take effect.

**2006:** Low sulfur diesel fuel required nationwide.

**2007:** Tighter emission standards for heavy-duty diesel trucks take effect nationwide.

**2009:** Greenhouse gas emission standards for passenger cars and light trucks.

**2011:** Tighter emission standards for off-road diesel equipment.

---

## **Web Resources** (*www.arb.ca.gov/californiaalmanac*)

Much of the information used to develop the Almanac is accessible through a variety of databases and tools available on the ARB website at *www.arb.ca.gov/californiaalmanac*.

### **Data**

**Real-time Air Quality Data** - Air Quality and Meteorological Information System (AQMIS2) - Allows access to near real-time air quality and meteorological data. These data are available in tabular summary reports.

**Historical Air Quality Data** - Aerometric Data Analysis and Management System (iADAM) - Allows access to historical data (data for record) in tabular summary reports or displayed as graphs.

**Emission Inventory Data** - Allows access to historical and projected emissions, vehicle activity, and human population. Data is available for 2004, as well as for the years 1975-2020 at five year intervals.

**Facility Search Engine** - Allows users to locate criteria or toxics emissions data for a specific facility.

**Top 25 Source Categories** - Provides users with emissions for the top 25 highest emitting source categories by geographic area.

**Community Level Emissions** - Community Health Air Pollution Information System (CHAPIS) - Allows users to query and view emissions using a map interface.

**Toxics Emission Factors** - California Air Toxics Emission Factor database (CATEF) - Provides over 2000 emission factors to estimate toxic air emissions for specific industrial processes or emissions.

### **Information**

**Area Designations** - Provides information regarding the designation areas in California with respect to the State Ambient Air Quality Standards.

**Biomass Burning Alternatives** - Information on alternatives to burning and provides helpful information on service providers in this effort.

**Central California Air Quality Studies (CCAQS)** - Comprises two studies with the goal of providing an improved understanding of particulate matter and visibility in central California.

**Outdoor residential waste burning information** - Provides information on regulations for outdoor residential waste burning and includes resources on alternatives to burning.

**Air Quality Data Quality Assurance Site Index** - Air monitoring web site that allows the user to gain access to the most recent quality assurance information on any particular air monitoring site. This information consists of pollutants monitored, location, operation information, and photos of the site if available.

**Transport** - Information on the movement of ozone and ozone precursors between basins or regions and established mitigation requirements.

## *Glossary of Air Quality Terms*

**Air:** So called “pure” air is a mixture of gases containing about 78 percent nitrogen; 21 percent oxygen; less than one percent of carbon dioxide, argon, and other gases; and varying amounts of water vapor.

**Air Basin:** A land area with generally similar meteorological and geographic conditions throughout. To the extent possible, air basin boundaries are defined along political boundary lines and include both the source and receptor areas. California is currently divided into 15 air basins.

**Air District:** A political body responsible for managing air quality on a regional or county basis. California is currently divided into 35 air districts.

**Air Monitoring:** Sampling for and measuring of pollutants present in the atmosphere.

**Air Pollution:** Degradation of air quality resulting from unwanted chemicals or other materials occurring in the air.

**Air Pollution Control District (APCD):** A county agency with authority to regulate stationary, indirect, and area sources of air pollution (e.g., power plants, highway construction, and housing developments) within a given county, and governed by a district air pollution control board composed of the elected county supervisors.

**Air Quality Management District (AQMD):** A group of counties or portions of counties, or an individual county specified in law with authority to regulate stationary, indirect, and area sources of air pollution within the region and governed by a regional air pollution control board comprised mostly of elected officials from within the region.

**Air Quality Management Plan (AQMP):** A plan prepared by an APCD / AQMD, for a county or region designated as a nonattainment area, for the purpose of bringing the area into compliance with the requirements of the national and/or California ambient air quality standards. AQMPs are incorporated into the State Implementation Plan (SIP).

**Air Quality Standard (AQS):** The prescribed level of a pollutant in the outside air that should not be exceeded during a specific time period to protect public health. Established by both federal and state governments.

**Air Toxics:** A generic term referring to a harmful chemical or group of chemicals in the air. Substances that are especially harmful to health, such as those considered under U.S. EPA’s hazardous air pollutant program or California’s AB 1807 and / or AB 2588 air toxics programs, are considered to be air toxics. Technically, any compound that is in the air and has the potential to produce adverse health effects is an air toxic.

**Ambient Air Quality Standards (California-CAAQS or National-NAAQS):** Health- and welfare-based standards for outdoor air which identify the maximum acceptable average concentrations of air pollutants during a specified period of time.

**Area-wide Sources (also known as “Area Sources”):** Stationary sources of pollution (e.g., water heaters, gas furnaces, fireplaces, and woodstoves) that are typically associated with homes and non-industrial sources. Area-wide sources do not include mobile sources. The California Clean Air Act requires air districts to include area-wide sources in the development and implementation of their Air Quality

Maintenance Plan. Under the federal air toxics program, an area-wide source is defined as any source that emits less than 10 tons per year of a single hazardous air pollutant (HAP) or 25 tons per year of all HAPs.

**Attainment Area:** A geographical area identified to have air quality as good as, or better than, the national and/or California ambient air quality standards. An area may be an attainment area for one pollutant and a nonattainment area for others.

**California Clean Air Act (CCAA):** A California law passed in 1988 which provides the basis for air quality planning and regulation independent of federal regulations. A major element of the Act is the requirement that local air districts in violation of the CAAQS must prepare attainment plans which identify air quality problems, causes, trends, and actions to be taken to attain and maintain California's air quality standards by the earliest practicable date.

**Criteria Air Pollutant:** An air pollutant for which acceptable levels of exposure can be determined and for which an ambient air quality standard has been set. Examples include: ozone, carbon monoxide, nitrogen dioxide, sulfur dioxide, and PM10 and PM2.5.

**Climate Change:** An change in the temperature of the Earth's troposphere. Climate change has occurred in the past as a result of natural influences, but the term is most often used to refer to the warming predicted by computer models to occur as a result of increased emissions of greenhouse gases.

**Emission Inventory:** An estimate of the amount of pollutants emitted into the atmosphere from major mobile, stationary, area-wide, and natural source categories over a specific period of time such as a day or a year.

**Emission Standard:** The maximum amount of a pollutant that is allowed to be discharged from a polluting source such as an automobile or smoke stack

**Environmental Justice:** The fair treatment of people of all races and incomes with respect to development, implementation, and enforcement of environmental laws, regulations, and policies.

**Expected Peak Day Concentration (EPDC):** See Peak 1-Hour Indicator

**Exceedance:** A measured level of an air pollutant higher than the national or state ambient air quality standards.

**Exposure:** The concentration of the pollutant in the air multiplied by the population exposed to that concentration over a specified time period.

**Exposure Assessment:** Measurement or estimation of the magnitude, frequency, duration and route of exposure to a substance for the populations of interest.

**Federal Clean Air Act (FCAA):** A federal law passed in 1970 and amended in 1974, 1977 and 1990 which forms the basis for the national air pollution control effort. Basic elements of the act include national ambient air quality standards for major air pollutants, mobile and stationary control measures, air toxics standards, acid rain control measures, and enforcement provisions.

**Hydrocarbon:** A general term used to describe compounds comprised of hydrogen and carbon atoms. Hydrocarbons are classified as to how photochemically reactive they are: relatively reactive or relatively non-reactive.

**Mean:** Average.

**Mobile Sources:** Sources of air pollution such as automobiles, motorcycles, trucks, off-road vehicles, boats, and airplanes (compare with Stationary Sources).

**Nonattainment Area:** A geographic area that does not meet either a State or federal standard for a given pollutant. This area usually consists of an air basin, but can be any geographic area defined by either the U.S. EPA or CARB.

**Nonattainment Transitional:** A subcategory of the nonattainment designation category for State standards that signals progress and implies the area is nearing attainment.

**Peak 1-Hour Indicator:** Using a statistical process, it is a site-specific and pollutant-specific value that represents the concentration expected to be exceeded once per year, on average, based on the distribution of data for the monitoring site. The calculation procedure uses data collected at the monitoring site for a three-year period. For example, the 2003 peak indicator is calculated using data for the years 2001, 2002, and 2003. The site with the highest peak indicator for a region is used for the long-term trends in the almanac. It is also referred to as the California Design Value or the Expected Peak Day Concentration.

**Precursor Emissions:** Emissions which form pollutants in the atmosphere due to the reaction of themselves with each other or with sunlight. Ozone is formed in the atmosphere when hydrocarbon and  $\text{NO}_x$  react in the presence of sunlight. Particulate Matter (PM) is a complex pollutant that can be formed from the reaction of gaseous precursors such as  $\text{NO}_x$ , ROG,  $\text{SO}_x$ , and ammonia.

**Reactive Organic Gas (ROG):** A reactive chemical gas, composed of non-methane hydrocarbons that may contribute to the formation of smog. Also sometimes referred to as non-methane organic gases (NMOGs).

**Risk Assessment:** An evaluation of risk which estimates the relationship between exposure to a harmful substance and the likelihood that harm will result from that exposure.

**State Implementation Plan (SIP):** A plan prepared by states and submitted to U.S. EPA describing how each area will attain and maintain national ambient air quality standards. SIPs include the technical foundation for understanding the air quality (e.g., emission inventories and air quality monitoring), control measures and strategies, and modeling analyses.

**Stationary Sources:** Non-mobile sources such as power plants, refineries, and manufacturing facilities which emit air pollutants (compare with Mobile Sources).

**Total Organic Gases (TOG):** All gases consisting of substances containing carbon, except carbon monoxide, carbon dioxide, carbonic acid, metallic carbides or carbonates, and ammonium carbonate.

**Toxic Air Contaminant (TAC):** An air pollutant, identified in regulation by the ARB, which may cause or contribute to an increase in deaths or in serious illness, or which may pose a present or potential hazard to human health. TACs are considered under a different regulatory process (California Health and Safety Code section 39650 et seq.) than pollutants subject to CAAQSs. Health effects to TACs may occur at extremely low levels, and it is typically difficult to identify levels of exposure which do not produce adverse health effects.

**Vehicle Miles Traveled (VMT):** The miles traveled by motor vehicles over a specified length of time (e.g., daily, monthly, or yearly) or over a specified road or transportation corridor.

**Volatile Organic Compounds (VOC):** A group of chemicals that react in the ambient air with nitrogen oxides and hydrocarbons in the presence of heat and sunlight to form ozone. Examples of VOCs include gasoline fumes and oil-based paints. This group of chemicals does not include methane or other compounds determined by U.S. EPA to have negligible photochemical reactivity.