ARB CCS Technical Discussion Series: CO2 EOR

Briana Mordick
Senior Scientist
Natural Resources Defense Council

August 23, 2016

CO2-EOR Production & CO2 Supply

- In 2012, CO2-EOR fields produced about 353,921 barrels of oil per day (bbl/d) in the U.S., which amounts to roughly 5.5% of domestic crude oil production. [1]
- Additional CO2-EOR production is limited by scarcity of CO2 supply
- ~ 80% of the CO2 used for EOR comes from naturally occurring, underground accumulations of CO2 [2]
- Remaining ~20% comes primarily from natural gas processing and fertilizer production plants [2]
- CO2 demand from natural sources will most likely continue to exceed supply, creating an incentive for CO2-EOR operators to pursue opportunities to capture man-made CO2 from large industrial facilities.

CO2-EOR Potential

	Technically Recoverable Oil (Billion Barrels)		Economically Recoverable Oil (Billion Barrels)		Economic CO2 Demand/Storage (Million Metric Tons)	
Oil price / CO2 price	State of Art	Next Generation	State of Art	Next Generation	State of Art	Next Generation
$85/bbl / $40/tCO2	61.4	120.3	26.9	67.2	10,430	19,930
$45/bbl / $23.25/tCO2	61.4	120.3	13.9	49.3	4,800	13,400

Source [3]

Sequestration Potential

- Total benefit from sequestering CO2 through EOR equivalent to ~1-3.5 times annual U.S. energy-related CO2 emissions, depending on oil & CO2 price and technology used. [3]



Source [4]

CO2-EOR Challenges

- Fields with long operating histories and large numbers of wells
- Observed quality of construction and maintenance in the field varies considerably with location, operator and age
- May have multiple owners over time and well files and well histories may be missing or incomplete
- State and federal records may also be incomplete, given that oil exploration and production in the U.S. predated regulation by several decades in some cases
- Location and plugging status of all wells within the field may be unknown or imprecisely known
- Orphan wells are ubiquitous in regions that have undergone oil and gas exploration and are a potentially significant leakage risk for CO2-EOR projects
- Orphan wells may have been constructed and/or plugged using outdated methods, and may have not been maintained over time to ensure that the integrity of the construction and plugging materials has not degraded

CO2-EOR Challenges

- Maintaining mechanical integrity critical to preventing leakage; performance of existing projects generally not well documented/publicly available
- Numerous studies demonstrate that determining the precise number of violations, mechanical integrity and contamination incidents, and their variation with time or location is not possible with existing data [5]
- Information is incomplete, outdated, or nonexistent making it is difficult to infer exact mechanical integrity failure rates or the number, extent and frequency of contamination or leakage incidents
- Class II regulations leave significant discretion to operators
- 40 states have primacy for Class II, contain 94% of Class II wells and produced 99% of U.S. onshore oil in 2015
- Significant variation exists among state rules in terms of stringency and completeness, as well as in enforcement

CO2-EOR Challenges

- The commercial time horizon for a CO2-EOR flood (a few years to decades) is shorter than the time horizon of interest for achieving effective sequestration of CO2 from the atmosphere (centuries or longer)
- Migration of CO2 out of pattern, into authorized zones, or to the atmosphere is possible after injection and production cease due to lack of focus on post injection site closure
- Change in ownership may result in inconsistent stewardship of the stored CO2 and/or loss of critical knowledge of the field and its operating history
- Development of advanced EOR techniques could result in operators reentering CO2-EOR fields at a future date to recover additional reserves, which could necessitate “blowing down” the field

Leakage Pathways and Rates

- Largest documented release rate occurred at a blowout at the Sheep Mountain CO2 Dome in Colorado.
  - Estimated 7,000 to 11,000 tons of CO2 were released per day for 17 days, resulting in a total of approximately 200,000 tons of leaked CO2. [6]
- Modeled CO2 leakage rates from a completely unobstructed pipe found a maximum hypothetical CO2 flow rate of approximately 20,000 metric tons/day (assuming a 7” inside diameter well and depth of ~5,000 feet)
  - Maximum exit gas velocity and flow rate is limited by the speed of sound
  - This theoretical maximum rate is approximately two times greater than Sheep Mountain [7]
- Maximum estimated leak rate from Aliso Canyon = 1,392 metric tons/day
- Leakage can occur through combination of well and geologic pathways, e.g. Leroy and Yaggy underground gas storage facilities [8][9]

Solutions

- Detailed site characterization to determine geologic suitability
- Site-specific Area of Review and Corrective Action plans that take into account field data and operating history
- Robust methods for identifying existing wells, including
  1. Historical Record Review
  2. Site Reconnaissance
  3. Aerial and Satellite Imagery Review, and;
  4. Geophysical and Air Emissions Surveys
- Ensuring MI of existing wells, including
  1. Well Record Review
  2. Field Inspection and Testing, and;
  3. Corrective Action
- Best practices for new well construction and conversion of existing wells
- Robust leak inspection, detection, reporting, and repair standards
- Comprehensive Mechanical Integrity Testing Plan, including post-closure

NRDC

References

- [1] Data for miscible and immiscible CO2 floods from: Oil & Gas Journal. (2012). Producing CO2, Other Gas and Chemical EOR in the US. Retrieved July 6, 2016 from Oil & Gas Journal website: <http://www.ogj.com/content/dam/ogj/print-articles/Volume%20110/April%202/Producing%20CO2%20Other%20Gas.pdf>; Data for total U.S. domestic production in 2012 of 6,482,000 bbl/d from Energy Information Administration. (2016). "Crude Oil Production." Crude Oil Production. Energy Information Administration. Retrieved July 6, 2016 from Energy Information Administration website: <http://www.eia.gov/dnav/pet/pet_crd_crpdn_adc_mbblpd_a.htm>.

- [2] Murrell, G., & DiPietro, P. (2013). North American CO2 Supply and Developments. In 19th Annual CO2 Flooding Conference

- [3] Kuuskraa, V. A., Van Leeuwen, T., Wallace, M., & DiPietro, P. (2011). Improving Domestic Energy Security and Lowering CO2 Emissions with "Next Generation" CO2-Enhanced Oil Recovery (CO2-EOR). National Energy Technology Laboratory, Pittsburgh, PA, USA. and 2016 unpublished custom analysis by Advanced Resources International for NRDC

- [4] U.S. Environmental Protection Agency. (2016). Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990 – 2014. Washington, D.C.: U.S. Environmental Protection Agency. EPA 430-R-16-002.

- [5] See, e.g. Koplos, J., Kobelski, B, Karimjee, A. and Sham C. H. (2006). UIC Program Mechanical Integrity Testing: Lessons for Carbon Capture and Storage?. Fifth Annual Conference on Carbon Capture and Sequestration - DOE/NETL, May 8 - 11, 2006.; Porse, S. L., Wade, S., & Hovorka, S. D. (2014). Can We Treat CO2 Well Blowouts like Routine Plumbing Problems? A Study of the Incidence, Impact, and Perception of Loss of Well Control. Energy Procedia, 63, 7149-7161.; U.S. Government Accountability Office. (2016). Drinking Water, EPA Needs to Collect Information and Consistently Conduct Activities to Protect Underground Sources of Drinking Water. Washington, D.C.: U.S. Government Accountability Office.; Watson, T., & Bachu, S. (2009). Evaluation of the Potential for Gas and CO2 Leakage along Wellbores. SPE Drilling & Completion, 24(1), 115-126.; Dusseault, M.B., Jackson, R.E., & MacDonald, D. (2014). Towards a Road Map for Mitigating the Rates and Occurrences of Long-Term Wellbore Leakage. Geofirma Engineering Ltd.

- [6] Wilson, E. J., Friedmann, S. J., & Pollak, M. F. (2007). Research for Deployment: Incorporating Risk, Regulation, and Liability for Carbon Capture and Sequestration. Environmental Science & Technology, 41(17), 5945-5952.

- [7] Aines, R. D., Leach, M. J., Weisgraber, T. H., Simpson, M. D., Julio Friedmann, S., & Bruton, C. J. (2009). "Quantifying the potential exposure hazard due to energetic releases of CO2 from a failed sequestration well". Energy Procedia, 1(1), 2421-2429.

- [8] Lewicki, J. L., Birkholzer, J., & Tsang, C. F. (2006). Natural and industrial analogues for release of CO2 from storage reservoirs: Identification of features, events, and processes and lessons learned.

- [9] Allison, M. L. (2001). Hutchinson, Kansas: A geologic detective story. Geotimes, 46(10).