



# Volatile organic compound emissions from green waste composting: Characterization and ozone formation

Anuj Kumar<sup>a, 1</sup> ✉, Christopher P. Alaimo<sup>a</sup>, Robert Horowitz<sup>b</sup>, Frank M. Mitloehner<sup>c</sup>, Michael J. Kleeman<sup>a</sup>, Peter G. Green<sup>a</sup> ✉

Show more ▾

☰ Outline | 🔗 Share 🗒 Cite

<https://doi.org/10.1016/j.atmosenv.2011.01.014>

[Get rights and content](#)

## Abstract

Composting of green waste separated from the disposed solid waste stream reduces biodegradable inputs into landfills, and contributes valuable soil amendments to agriculture. Agencies in regions with severe air quality challenges, such as California's San Joaquin Valley (SJV), have raised concerns about gases emitted during the composting process, which are suspected to contribute to persistent high levels of ground-level ozone formation. The goal of the current study is to thoroughly characterize volatile organic compound (VOC) emissions from green waste compost piles of different ages (fresh tipped piles, 3–6 day old windrows, and 2–3 week old windrows). Multiple sampling and analytical approaches were applied to ensure the detection of most gaseous organic components emitted. More than 100 VOCs were detected and quantified in this study, including aliphatic alkanes, alkenes, aromatic hydrocarbons, biogenic organics, aldehydes, ketones, alcohols, furans, acids, esters, ether, halogenated hydrocarbons and dimethyl disulfide (DMDS). Alcohols were found to be the dominating VOC in the emissions from a compost pile regardless of age, with fluxes ranging from 2.6 to 13.0 mg m<sup>-2</sup> min<sup>-1</sup> with the highest emissions coming from the younger composting windrows (3–6 days). Average VOC emissions other than alcohols were determined to be 2.3 mg m<sup>-2</sup> min<sup>-1</sup> from younger windows, which was roughly two times higher than either the fresh tipping pile (1.2 mg m<sup>-2</sup> min<sup>-1</sup>) or the older windrows (1.4 mg m<sup>-2</sup> min<sup>-1</sup>). It was also observed that the older windrows emit a slightly larger proportion of more reactive compounds. Approximately 90% of the total VOCs were found

to have maximum incremental reactivity of less than 2. Net ozone formation potential of the emissions was also assessed.

## Highlights

► Characterization of VOC emissions from green waste compost of different ages. ► Emissions are dominated by small alcohols ranging from 66–85% of the total. ► Young windrows (3–6 days) had the highest flux, but less reactivity to form ozone. ► Older compost windrows (2–3 weeks) exhibited lower fluxes, but more reactivity. ► Field ozone assays and model calculations confirm low ozone formation potential.

 **Previous**

**Next** 

## Keywords

Green waste; Compost; VOC; Organic matter; Chemical emissions

---

[Recommended articles](#)

---

## Cited by (59)

[Odorous gas emissions from sewage sludge composting windrows affected by the turning operation and associated health risks](#)

2022, Science of the Total Environment

[Show abstract](#) ✓

[Measuring volatile emissions from biosolids: A critical review on sampling methods](#)

2022, Journal of Environmental Management

[Show abstract](#) ✓

[Reduction in VOC emissions by intermittent aeration in bioreactor landfills with gas-water joint regulation](#)

2021, Environmental Pollution

[Show abstract](#) ✓

## Performance evaluation of a full-scale open bed biofilter through on-site measurements and CFD analyses

2021, Process Safety and Environmental Protection

[Show abstract](#) ✓

## First-principles insight on interaction behavior of diethylbenzene and ethyltoluene on $\alpha$ -arsenene nanoring

2020, Materials Today Communications

[Show abstract](#) ✓

## Characteristics of atmospheric volatile organic compounds in urban area of Beijing: Variations, photochemical reactivity and source apportionment

2020, Journal of Environmental Sciences (China)

[Show abstract](#) ✓



[View all citing articles on Scopus](#)

<sup>1</sup> Current address: Carlsbad Environmental Monitoring & Research Center, New Mexico State University, 1400 University Drive, Carlsbad, NM 88220, USA.

[View full text](#)

Copyright © 2011 Elsevier Ltd. All rights reserved.



Copyright © 2022 Elsevier B.V. or its licensors or contributors.  
ScienceDirect® is a registered trademark of Elsevier B.V.

