

DRAFT COPY

Chemical Consequences of Air Quality Standards and of Control Implementation
Programs: Roles of Hydrocarbons, Oxides of Nitrogen, Oxides of Sulfur and
Aged Smog in the Production of Photochemical Oxidant and Aerosol

Contract No. 5-385
California Air Resources Board
Final Report
May 1977

Principal Investigator

Dr. James N. Pitts, Jr.

Program Manager

Dr. Arthur M. Winer

Program Research Staff

Dr. Karen R. Darnall
Dr. George J. Doyle
Dr. Richard A. Graham
Dr. Ernesto C. Tuazon

Technical Support Staff

Mr. Frank R. Burleson
Mr. Dennis R. Fitz
Mr. L. Michael Kienitz
Mr. Glen C. Vogelaar
Ms. Laurie A. Willis

Contributing Research Staff

Dr. William P. Carter
Dr. William C. Kuby
Dr. Alan C. Lloyd
Ms. Minn P. Poe
Dr. Edgar R. Stephens

TABLE OF CONTENTS

	<u>Page</u>
Abstract	ii
List of Figures	iii
List of Tables	iv
Executive Summary	v
Acknowledgments	xiii
I. Introduction	1
II. Facilities and Methods	3
A. Chamber Facility and Analytical Methods	3
B. Experimental Procedures	7
III. Development of a Reactivity Scale and Classification for Organics Based on Their Rates of Reaction with the Hydroxyl Radical	11
A. Relative Rates of Reaction of Organic Compounds with the Hydroxyl Radical	11
B. Development of a Reactivity Classification Based on the Reaction of Organics with OH	14
IV. A Chamber Study of Sulfate Aerosol Formation	32
V. Further Development of an Air Parcel Transport Simulation Methodology	39
VI. Ambient Air Analysis	47
VII. References	48
VIII. List of Publications from SAPRC-ARB Chamber Program	56
Appendix A. Reprints of OH Rate Constant Determination and Reactivity Scale Papers	58
Appendix B. Inorganic and Hydrocarbon Data for AGC Runs 216 through 223	101
Appendix C. Plots of $\ln SO_2$ vs. time for AGC Runs 216 through 222	148

ABSTRACT

The environmental chamber facility established at the Statewide Air Pollution Research Center, University of California, Riverside, under a joint Air Resources Board--University of California (Project Clean Air) program has been employed in investigations:

To determine the relative rates of reaction of the hydroxyl (OH) radical with a large number of organic compounds including those representative of (a) combustion products, (b) solvent emissions, (c) natural emissions and (d) potential aerosol precursors.

To develop, in conjunction with the above experimental work, a new reactivity scale and classification of organic compounds based on their rates of reaction with the OH radical.

To provide data concerning the photochemically-promoted conversion of sulfur dioxide (SO_2) to sulfate aerosol.

Previously developed procedures for characterizing polluted ambient air parcels utilizing a diffusion model and available meteorological and air monitoring data have been further refined:

To determine the rates of dilution and diffusion for various pollutants in characteristic air parcels and to utilize this information to design chamber experiments simulating transport, new pollutant emissions and dilution phenomenon.

This report was submitted in fulfillment of contract No. 5-385 by the Statewide Air Pollution Research Center, University of California, Riverside, under the partial sponsorship of the California Air Resources Board. Work was completed as of January 31, 1977.

LIST OF FIGURES

<u>Figure No.</u>		<u>Page</u>
1	Schematic Drawing of Glass Chamber Facility	4
2	Air Purification System	8
3	Comparison of Relative Rates of Hydrocarbon Disappearance Determined by the Chamber Method with Selected Published Rate Constants for Reaction of Those Hydrocarbons with OH Radicals	12
4	Predicted Relative Importance of Several Reactive Intermediates during Photooxidation of a Propene-n-Butane Mixture under Simulated Atmospheric Conditions	13
5	Concentrations of Selected Organic Compounds during Irradiation of a HC-NO _x Mixture in Air at 305 ± 2 K	16
6	Concentrations of Ketones and Isobutene during Irradiation of a HC-NO _x Mixture in Air at 305 ± 2 K	17
7	Concentrations of Ethers and Isobutene during Irradiation of a HC-NO _x Mixture in Air at 305 ± 2 K	18
8	Size Distributions (Whitby) of Spray-Dried Potassium Sulfate Solutions in SAPRC All-Glass Chamber	38
9	Volume of Chamber Sample to be Removed Each Hour (VDIL) for Simulation of NO _x Concentration in Ambient Air Parcels	41
10	Volume of Pure NO _x to be Added to Chamber Each Hour (VIN) for Simulation of NO _x Concentration in Ambient Air Parcels	42
11	Volume of Chamber Sample to be Removed Each Hour (VDIL) for Simulation of Hydrocarbon Concentration in Ambient Air Parcels	43
12	Volume of Pure Hydrocarbon to be Added to Chamber Each Hour (VIN) for Simulation of Hydrocarbon Concentration in Ambient Air Parcels	44
13	Hourly Average Concentrations of NO _x and Hydrocarbons for Ambient Air Parcel	46

LIST OF TABLES

<u>Table No.</u>		<u>Page</u>
1	Rate Constants for Reaction of OH with Organic Compounds Obtained Using the Chamber Method	vii
2	Proposed Reactivity Classification of Organics (and Selected Inorganics) Based on Reaction with the Hydroxyl Radical	ix
3	Rate of SO ₂ Loss in All-Glass Chamber under a Variety of Conditions	xi
4	Chemical and Physical Parameters Measured in Glass Chamber Studies	5
5	Rate Constants for Reaction of OH with Organic Compounds Obtained Using the Chamber Method	15
6	Rate Constants for Reaction of OH with Organic Compounds Obtained Using the Chamber Method	19
7	Reactivity Scale for Organics Based on Rate of Consumption Due to Reaction with the Hydroxyl Radical	21
8	Rate constants for Reaction of Organics (and Selected Inorganics) with the Hydroxyl Radical and the Corresponding Reactivity Classification	22
9	Proposed Reactivity Classification of Organics (and Selected Inorganics) Based on Reaction with the Hydroxyl Radical	28
10	California Air Resources Board (ARB) Reactivity Classification of Organic Compounds	30
11	Rate of SO ₂ Loss in All-Glass Chamber under a Variety of Conditions	34
12	Aerosol Measurements Using the Whitby and Climet Analyzers during Propene-Ozone-SO ₂ Experiment	35

EXECUTIVE SUMMARY

Despite two decades of intensive research concerning the photochemistry of polluted atmospheres, and the concerted attention during the past ten years of local, state and federal air pollution control agencies to the problem of reducing photochemical oxidant in urban areas, substantial uncertainty still remains as to the most cost-effective and societally acceptable means of solving this major environmental problem. In order to provide an experimental data base relating to the major unresolved issues of photochemical smog formation, the SAPRC environmental chamber facility was designed and constructed (1970-73) with support from the California Air Resources Board. The facility was initially (1973-75) utilized in a study of oxidant formation from hydrocarbon-oxides of nitrogen (HC-NO_x) mixtures irradiated under conditions simulating those found in the California South Coast Air Basin. Using a twelve-hydrocarbon mixture as a surrogate for the ambient pollutant mix, HC and NO_x concentrations ranging from those found in present-day polluted atmospheres down to those to be expected from the implementation of emission control strategies aimed at meeting the State and Federal ambient air quality standards were investigated. In conjunction with this program evaluations of the accuracy or specificity of the analytical methods for two major pollutants (ozone and oxides of nitrogen) were also carried out.

In response to some of the most recent developments in atmospheric chemistry and pollution control, the original precursor-secondary pollutant study has been extended or applied in the current year to

- determination of OH radical rate constants for a large number of organic compounds under simulated atmospheric conditions in an environmental chamber
- development of a new reactivity scale based on the reaction of organics with the OH radical
- a chamber study of the loss of SO₂ and formation of sulfate aerosol under simulated atmospheric conditions, and
- development of methods to more accurately simulate in smog chamber experiments the ambient conditions of transport, input of new emissions, and dilution.

OH Radical Rate Constants. The use of relative rates of reactions of hydrocarbons determined in the SAPRC all-glass environmental chamber under

simulated atmospheric conditions was shown two years ago to be a reliable method for determining the OH rate constants for these compounds. The validation of the chamber method for OH radical rate constant determinations was based on: (a) the good to excellent agreement between rate constants determined using this method and those determined by elementary methods with the individual hydrocarbons (b) kinetic computer models of photochemical smog formation which uniformly indicate the OH radical to be the major reactive intermediate involved in consumption of hydrocarbons (other than alkenes which also react appreciably with ozone) and (c) support of the model predictions by the direct and indirect observation of OH radical concentrations in both ambient air and experimental systems at about the expected levels of 10^6 radicals cm^{-3} .

Utilizing the chamber method we have now determined OH rate constants for 32 additional organic compounds falling into four classes of potential importance in the production of photochemical air pollution (see Table 1) (i.e., combustion source organics, solvent components, naturally occurring hydrocarbons and potential aerosol precursors). The relative rates of disappearance which were obtained from the chamber experiments were converted to absolute rate constants using either n-butane or isobutene as a reference compound having a well established OH rate constant.

Some advantages of the chamber method include the ability to determine rates for several compounds in each series of experiments, the low purity requirements for the compounds studied (as long as interferences do not occur in the gas chromatographic analyses) and the ability to study compounds not amenable to elementary methods.

Reactivity Scale Based on Reaction with The OH Radical. We have proposed a hydrocarbon reactivity scale based on the primary chemical act of hydroxyl radical (OH) attack on organic species. As discussed above, the OH radical is the key reactive intermediate in photochemical air pollution. Moreover, a reactivity scale based on the depletion of hydrocarbons by reactions with OH has utility in assessing hydrocarbon chemical behavior in polluted ambient air, since only those compounds which participate at "significant" rates in atmospheric reactions are of consequence in the chemical transformations in ambient air.

Table 1. Rate Constants for Reaction of OH with Organic Compounds
Obtained Using the Chamber Method

Compound	Rate Constant 10^{-9} (ℓ mole $^{-1}$ sec $^{-1}$)	Reference
Combustion Sources		
Ethene	5.2 ± 1.0	a
Propene	17.5 ± 3.5	a
cis-2-Butene	14.9 ± 2.2	b
	39.2 ± 8.0	a
	37.9 ± 5.6	b
	32.9 ± 6.6	c
1,3-Butadiene	46.4 ± 9.3	a
2,3-Dimethylbutane	3.1 ± 0.5	f
Isopentane	2.0 ± 0.4	a
2-Methylpentane	3.2 ± 0.6	a
3-Methylpentane	4.3 ± 0.9	a
n-Hexane	3.8 ± 0.8	a
m-Xylene	12.9 ± 2.6	a
n-Propylbenzene	3.7 ± 0.8	a
Isopropylbenzene	3.7 ± 0.8	a
Ethylbenzene	4.8 ± 1.0	a
o-Ethyltoluene	8.2 ± 1.6	a
m-Ethyltoluene	11.7 ± 2.3	a
p-Ethyltoluene	7.8 ± 1.6	a
Solvent Components		
Methyl ethyl ketone	2.0 ± 0.6	b
Methyl isobutyl ketone	9.0 ± 3.0	b
Diisobutyl ketone	15 ± 5	b
Isopropyl alcohol	4.3 ± 1.3	d
Diethyl ether	5.6 ± 1.1	d
Di-n-propyl ether	10.4 ± 2.1	d
Tetrahydrofuran	8.8 ± 1.8	e
n-Propyl acetate	2.6 ± 0.5	e
sec-Butyl acetate	3.4 ± 0.7	e
Naturally Occurring Hydrocarbons		
α -Pinene	34.8 ± 5.2	b
β -Pinene	40.6 ± 6.1	b
d-Limonene	90.0 ± 13.5	b
Potential Aerosol Precursors		
2,2,3-Trimethylbutane	2.3 ± 0.5	f
1-Heptene	22 ± 5	f
Cyclohexene	47 ± 9	f
1-Methylcyclohexene	58 ± 12	f

^aLloyd, Darnall, Winer and Pitts, J. Phys. Chem., 80, 789 (1976).

^bWiner, Lloyd, Darnall and Pitts, J. Phys. Chem., 80, 1635 (1976).

^cDarnall, Winer, Lloyd and Pitts, unpublished results, 1976.

^dLloyd, Darnall, Winer and Pitts, Chem. Phys. Lett., 42, 205 (1976)

^eWiner, Lloyd, Darnall and Pitts, Chem. Phys. Lett., submitted (1977).

^fDarnall, Winer, Lloyd and Pitts, Chem. Phys. Lett., 44, 415 (1976).

For purposes of comparison with other reactivity scales, we have proposed a five-class scale which is based on order of magnitude differences in reactivity toward OH relative to that of methane (=1). Table 2 shows the five classes, the OH rate constant range appropriate to each class and the reactivity assignment for more than 110 compounds for which OH rate constants have been determined.

A comparison of this reactivity classification with the one based on ozone formation that was recently adopted by the ARB (which is very similar to the one under consideration by the EPA) shows them to be quite consistent with a few minor exceptions.

Advantages of the proposed OH reactivity scale include (a) its ready extension to additional compounds once their rate of reaction with OH is known, (b) its divisibility into as many classes as is desirable for the purposes of devising control priorities for organic emissions, and (c) the greater weight (than given previously) to alkanes and aromatics which over long irradiation periods will react to a significant extent. Disadvantages include (a) its limited applicability to compounds which undergo significant photodissociation in the atmosphere or which react at an appreciable rate with other atmospheric species, e.g., alkenes reacting with ozone, and (b) the uncertainties resulting from the lack of information concerning the identity and fates of subsequent products.

A Chamber Study of Sulfate Aerosol Formation. In polluted atmospheres, one of the major sources of sulfate aerosol appears to be the conversion of sulfur dioxide in conjunction with the nitrogen oxide-promoted photooxidation of organic compounds. In the South Coast Air Basin the sulfate burden has been increasing at a rate of about 10% per year. Although this may worsen if increased use of high sulfur fuels occurs, there is also the possibility of a counteracting effect resulting from decreased hydrocarbon and NO_x emission and therefore decreased conversion rates. To provide a quantitative experimental data base for a number of such future scenarios, we have extended our basic precursor-secondary pollutant program to include a study of SO_2 loss and aerosol formation.

The initial phase of this program has been completed. The feasibility of using the SAPRC all-glass environmental chamber has been established by study of the decay of SO_2 in pure air under both irradiated and non-irradiated

Table 2. Proposed Reactivity Classification of Organics (and Selected Inorganics) Based on Reaction with the Hydroxyl Radical

Class I ($\leq 4.8 \times 10^7$) ^a	Class II ($4.8 \times 10^7 - 4.8 \times 10^8$) ^a	Class III ($4.8 \times 10^8 - 4.8 \times 10^9$) ^a	Class IV ($4.8 \times 10^9 - 4.8 \times 10^{10}$) ^a	Class V ($\geq 4.8 \times 10^{10}$) ^a
Trifluoromethane	Trichloromethane	Neopentane	Ethylbenzene	2-Methyl-2-butene
1,1,2-Trichloro-1,2,2-trifluoroethane	Dichloromethane	Methanol	n-Octane	3-Carene
Tetrafluoromethane	Carbon monoxide	Methylacetylene	Diethyl ether	1-Methylcyclohexene
1,2-Dichloro-1,1,2,2-tetrafluoroethane	Ammonia	2,2,3,3-Tetramethylbutane	p-Xylene	β -Phellandrene
Chlorotrifluoromethane	Acetylene	Cyclobutane	p-Ethyltoluene	Carvomenthene
Dichlorodifluoromethane	1,2-Dichloroethane	Benzene	o-Ethyltoluene	d-Limonene
Fluorotrichloromethane	1,2-Dibromoethane	Isobutane	o-Xylene	2,3-Dimethyl-2-butene
1-Chloro-1,1-difluoroethane	Ethane	Propane	Tetrahydrofuran	Dihydromyrcene
Tetrachloromethane	1,1-Dichloroethane	n-Butane	Methyl isobutyl ketone	Myrcene
Difluoromethane	Chloroethane	Ethanol	p-Cymene	cis-Ocimene
Methane		Methyl ethyl ketone	Di-n-propyl ether	
2-Chloro-1,1,1-trifluoroethane		Isopentane	m-Ethyltoluene	
2-Chloro-1,1,1,2-tetrafluoroethane		1-Propanol	Methoxybenzene	
1,1,1-Trichloroethane		2,2,4-Trimethylpentane	Methylamine	
Fluoromethane		2,2,3-Trimethylbutane	m-Xylene	
Dichlorofluoromethane		n-Propyl acetate	1,2,3-Trimethylbenzene	
2,2-Dichloro-1,1,1-trifluoroethane		Allene	Diisobutyl ketone	
1,1-Difluoroethane		2,3-Dimethylbutane	Propene	
Chlorofluoromethane		Hydrogen sulfide	3,3-Dimethyl-1-butene	
Bromomethane		2-Methylpentane	1-Pentene	
Chloromethane		sec-Butyl acetate	1-Hexene	
Chlorodifluoromethane		Toluene	1,2,4-Trimethylbenzene	
		Cyclopentane	Methyl mercaptan	
		n-Propylbenzene	o-Cresol	
		Isopropylbenzene	1-Butene	
		n-Hexane	1-Heptene	
		n-Pentane	1,3,5-Trimethylbenzene	
		p-Menthane	Isobutene	
		1-Butanol	cis-2-Butene	
		Cyclohexane	2-Methyl-1-butene	
		2-Propanol	o-Pinene	
		4-Methyl-2-pentanol	cis-2-Pentene	
		3-Methylpentane	β -Pinene	
		Ethene	trans-2-Butene	
			Cyclohexene	
			1,3-Butadiene	
			Isoprene	

^aRange of values (in liter mole⁻¹ sec⁻¹) for the rate constant for reaction of the OH radical with the listed compounds.

conditions at various relative humidity levels. Decay rates in the presence of ozone, ozone + propene and SAPRC surrogate have also been determined. These results are summarized in Table 3.

A brief study of aerosol formation was conducted during one of the SO₂-ozone-propene experiments. Particle size distributions monitored as a function of time indicated (1) a rapid growth of small particles following the addition of propene to a SO₂-ozone mixture and (2) the increase in larger-sized particles was accompanied by a decrease in the smaller-sized particles.

An experiment to investigate the behavior of a test aerosol (spray-dried potassium sulfate) in the all-glass chamber indicated (a) that to obtain at least semiquantitative data on aerosol yield, the system should be seeded and (b) that measurement of b_{scat} in chamber experiments will not give a reliable indication of visibility reduction potential because the chamber wall losses leave the aerosol severely depleted in the high light scattering region (0.2-1.0 μm).

Air Parcel Transport Simulation. To provide a basis for designing realistic chamber experiments simulating air parcel transport effects during a full day irradiation, we have developed procedures for determining the rates of dilution and diffusion for the various pollutants in an air parcel. Thus for a given trajectory which was determined by interpolation of ground level meteorological data, the ambient air quality was determined by a similar interpolation procedure using the air quality data. Based on the results for the CO concentration history, a diffusion model was generated to allow for the transport of pollutants in the vertical direction. Both dilution and fresh pollutant injection rates required to approximate air parcel histories were then determined.

Recently, the choice of dilution rates of approximately 10% per hour by workers operating the large outdoor smog chambers in North Carolina (and by workers in a number of previous chamber studies explicitly employing dilution) has been challenged by Shair^{1,2} as being too low. Our initial data suggest that the situation is complex and that both groups of workers are correct to some extent. For example, based on NO_x, the peak dilution we have calculated is approximately 50%--far above those considered by previous chamber workers. However, for an

Table 3. Rate of SO₂ Loss in All-Glass Chamber under a Variety of Conditions (using FC-12 as Inert Tracer for Determination of Dilution Rate)

AGC Run No.	RH Range (%)	Amount of Water (torr)	Temp. Range (°C)	Light Intensity	SO ₂ Loss Rate (% hr ⁻¹)	Estimated Net ^a SO ₂ Loss Rate (% hr ⁻¹)
216	1-2.5	0.6-1.0	32.6-33.7	0%		0.4
217 ^b	11-24.5	5.1-10.3	34.4-36.6	100%		3
218 ^c	40-49	15.5-18.9	33.2-33.8	0%		0.4
218 ^d	31-55.5	14.0-22.2	33.0-36.0	100%		5
219 ^e	81.5-71.5	30.1-26.9	32.6-34.2	0%		5
220 ^f	41-40	15.2-15.3	32.7-32.8	0%		30
220 ^g	40-38.5	14.9-14.5	32.8-33.1	0%		0.2
221 ^h	44-38.5	16.4-13.9	32.7-33.2	0%		1 ⁱ
222 ^j	46.3-32.5	15.9-12.7	34.3-30.5	0%		2
222 ^k	32.5-26.0	13.2-12.0	33.8-36.6	100%		4

^aTotal rate of SO₂ disappearance minus dilution rate (as determined from monitoring of FC-12).
^bContinuation of 216 at 100% light intensity; additional SO₂ injected; approx. 0.5 ppm ozone present in this experiment.

^cFlash lamp in Teco 43 replaced; instrument zeroed and spanned using MSA filter and calculated SO₂ in chamber, respectively.

^dOzone increased to ~0.3 ppm in first 6 hours of irradiation and remained at about that value.

^eLamp bases in Teco 43 replaced; zeroed and spanned as in b.

^f1 ppm propene and 0.6 ppm ozone present; average rate for first 30 min of reaction.

^g0.40 ppm propene and 0.20 ppm ozone present; average rate after 1.75 hr of reaction.

^h0.57 ppm ozone added.

ⁱValue for time after ozone was added.

^jSurrogate (~1700 ppbC) and NO_x (0.3 ppm calc.) present initially.

^kSurrogate (~1100 ppbC) and NO_x (~0.5 ppm) present at beginning of irradiation.

air parcel beginning at ground level, the average dilution rate over the entire trajectory, again based on NO_x , was 16%; and if one uses only that portion of the trajectory from 6:00 in the morning till 1:00 in the afternoon, the average dilution rate is 21%. The dilution rates based on hydrocarbon are considerably lower than the above values for NO_x . Thus, the results of this analysis for hydrocarbon appear to be more in line with the lower dilution values used in previous chamber studies.

The fact that dilution rates based on NO_x are considerably larger than those based on hydrocarbon is a crucial observation and results from the fact that the concentration of hydrocarbons is far in excess of the concentration of NO_x . Consequently, less volume need be removed from the chamber to achieve the mass fluxes that occur. Obviously, this situation presents a problem in determining an appropriate simulation in actual chamber experiments and work is continuing to determine what dilution volume is required to maintain an appropriate hydrocarbon-to- NO_x ratio, a possibly more realistic control parameter.

ACKNOWLEDGMENTS

Stimulating discussions and valuable exchanges of technical information, for which we express our appreciation, took place at various times during this program with the following members of the California Air Resources Board staff:

Mr. Frank Bonamassa

Dr. Robert C. Grant

Dr. John R. Holmes

Mr. Tom Quinn

Dr. Jack K. Suder

Dr. Leo Zafonte

We wish to specifically acknowledge the collaboration of Dr. Alan C. Lloyd in the hydroxyl radical rate constant studies and in development of the OH-reactivity scale, and of Dr. William C. Kuby in developing the methodology for designing air parcel transport simulation experiments. In addition, we thank our colleagues Dr. Roger Atkinson for many helpful discussions and Ms. Minn P. Poe for carrying out computer processing of the data.

Contributions from the Various Donors Fund of the Statewide Air Pollution Research Center, the U. S. Environmental Protection Agency (Grant No. R-800649), and the National Science Foundation--Research Applied to National Needs (Grant Nos. GI-41051 and AEN73-02904-A02), which supported in part the purchase of a computer data acquisition system employed in the ARB program, are gratefully acknowledged.

Finally, we thank the California Air Resources Board for their financial support of the work performed under this contract.

ALL OZONE AND OXIDANT DATA REPORTED IN THIS
STUDY WERE OBTAINED WITH INSTRUMENTS CALIBRATED
BY THE UB ABSORPTION METHOD ADOPTED BY THE
CALIFORNIA AIR RESOURCES BOARD, JUNE 1975.

The statements and conclusions in this report are those of the contractor and not necessarily those of the California Air Resources Board. The mention of commercial products, their source or their use in connection with material reported herein is not to be construed as either an actual or implied endorsement of such products.

I. INTRODUCTION

Events taking place in the area of control of photochemical air pollution, particularly in California, at the time the work presented here was proposed demonstrated anew the critical importance of approaching the air pollution problem on the strength of a valid technical, medical and socio-economic data base. Developments which illustrated this included (1) the strong public and legislative reaction to the Air Resources Board (ARB) NO_x retrofit program for 1966-1970 light duty vehicles in the South Coast Air Basin (SCAB), (2) the occurrence of discrepancies as large as factors of four in estimates of reactive hydrocarbons emitted in the SCAB by control agencies using differing reactivity scales,³ (3) the conversion of power plants in the SCAB from natural gas to fuel oil and the increasing use of high sulfur fuels, and (4) the proposals for relaxation of the implementation of all automobile emission standards as a trade off for improved fuel economy to help combat dependence on foreign oil.

In the face of these developments, some of which threaten to substantially slow the momentum of the early 1970's toward achieving significantly lower pollutant burdens, the following facts concerning air quality in the California South Coast Air Basin must be recognized:

- Pasadena exceeded the Federal air quality standard for oxidant of 0.08 ppm for 1 hour on 197 days in 1976.
- The state and Federal ambient air quality standards for nitrogen dioxide (NO₂) continue to be violated at many locations in the SCAB as widespread as Long Beach, Pasadena and Pomona.
- Atmospheric particulate nitrate which, according to a recent preliminary EPA study,⁴ may be associated with adverse health effects at levels estimated to be less than 10 micrograms per cubic meter, has been found at very high levels (about 70 micrograms per cubic meter, 24-hr average at one location) in the eastern portion of the Basin during the summer of 1975.
- Similarly, atmospheric particulate sulfate, which has been shown⁵ to be associated with adverse health effects at levels of 8-10 micrograms per cubic meter (24-hr average) has been observed at

levels as high as 20.4 micrograms per cubic meter (24-hr average) in West Covina.

In view of these developments and the apparent narrowing of control strategy options, the SAPRC environmental chamber laboratory has conducted a program designed to obtain quantitative data on chemical and physical transformations in polluted urban atmospheres of direct applicability to air pollution control strategies. In the current year research efforts were made in the following areas: (a) determination of OH radical rate constants under simulated atmospheric conditions in an environmental chamber for a large number of organic compounds, including those from combustion and other anthropogenic sources, as well as from natural sources, (b) development of a new reactivity scale based on the reaction of organics with the OH radical (c) a chamber study of the loss of SO₂ and formation of sulfate aerosol under simulated atmospheric conditions, and (d) development of methods to more accurately simulate in smog chamber experiments the ambient conditions of transport, input of new emissions, and dilution.

II. FACILITIES AND METHODS

A. Chamber Facility and Analytical Methods

The experiments in this study were carried out in a 6400-liter (226 ft³), all-glass (Pyrex) chamber^{6,7} which has a surface-to-volume ratio of 3.4 m⁻¹ (1.04⁻¹). The dimensions of the glass chamber are 8' x 8' x 4' (i.e., a flat box). Photolyzing radiation is provided by two externally mounted, diametrically opposed banks of forty Sylvania 40-W BL (black light) lamps, which are backed by arrays of Alzak-coated reflectors. The supporting analytical facilities employed in the glass chamber studies are shown schematically in Figure 1 and are described in detail below. The on-line computer data acquisition system which was installed in November 1974 and became fully operational in April, 1975, has been used throughout this program.

The current physical and chemical measurement methods and the range, precision, and accuracy of measurement of each of these parameters are given in Table 4. The details of the analytical and data processing procedures are described below for each species or variable monitored. The data obtained are computer processed and presented in tabular form for each run and have not been corrected for losses due to sampling from the chamber.

OZONE (O₃) was monitored by ultraviolet absorption analyzers (Dasibi-1003 or AH-1003). Until January 1976, these instruments were calibrated against 2% neutral buffered potassium iodide.⁸ Since it has been demonstrated^{9,10} that at ~50% RH this calibration procedure yields ozone values which are too high by a factor of ~1.25, all ozone data obtained prior to January 1976 have been corrected by a factor of 0.8. In January 1976 the instruments were respanned using UV absorption method adopted by the ARB in June, 1973.

NITROGEN OXIDES (NO, NO₂, and NO_x) were monitored by chemiluminescent detection (TECO 14B). The NO₂ and NO_x modes of this and similar chemiluminescent NO-NO_x analyzers have been shown to respond quantitatively to other nitrogen-containing compounds, such as peroxyacetyl nitrate (PAN) and organic nitrates and nitrites.^{11,12} All NO₂ and NO_x data reported




Figure 1. Schematic Drawing of Glass Chamber Facility.

Table 4. Chemical and Physical Parameters Measured in Glass Chamber Studies

Parameter	Range	Method	Sampling Rate	Precision	Accuracy
Ozone	0-20 ppm	UV Absorption Analyzer	570 ml/min	± 0.005 ppm	± 5%
NO	0-10 ppm	Chemiluminescent Analyzer	895 ml/min	± 2% F.S. By difference	± 5% By difference
NO ₂	0-10 ppm				
NO _x	0-10 ppm				
SO ₂	0-5 ppm	Pulsed Fluorescence	1000 ml/min	± 0.005 ppb	± 5%
PAN	>1 ppb	GC		± 0.5 ppb	± 10%
Formaldehyde	>10 ppb ^a	Chromotropic Acid	1 l/min (30 min samples)	± 5%	± 10%
Individual HCs	0.5-1 ppb 1-2 ppb >2 ppb	GC GC GC		± 15% ± 10% ± 2%	Limited by calibration standard (typically ± 3%) and/or by sampling techniques
CO	0-10 ppm	GC	100 ml/min	± 2% F.S.	± 10%
Temperature	15° to 40°C	Thermometer Thermocouple		± 0.02°C ± 0.1°F	± 0.5°C ± 2°F
Light Intensity	0-1 Solar Constant Relative	NO ₂ Actinometry Photodiode		± 5%	b
Relative Humidity	0-100%	Brady Array		± 1% RH	± 5%

^a Determined by practical values of sampling time and flow rate

^b See discussion on page 9.

here have been corrected by subtraction of measured or interpolated PAN concentrations.

SULFUR DIOXIDE (SO₂) was monitored by the pulsed fluorescence method (Teco 43). Approximate calibrations were obtained using chamber samples. A more precise calibration was carried out using a Monitor Labs model 8500 permeation type calibrator.

CARBON MONOXIDE (CO) was monitored by gas chromatography (Beckman-6800). The instrument was calibrated daily with a standard gas sample.

PEROXYACETYL NITRATE (PAN) was monitored by gas chromatography with electron capture detection (GC-ECD).^{13,14} Samples were taken in a 100-ml precision bore syringe and transferred to the GC sampling system as quickly as possible. Peak heights were read from the strip chart and converted to concentration units using a calibration function which was determined periodically.

FORMALDEHYDE (HCHO) was monitored using the chromotropic acid method.¹⁵ Air from the chamber was drawn through a bubbler at the rate of 1 l min⁻¹ and the total volume per sample was determined using a timer-controlled shutoff system. Generally, a 30-min. sample was taken. The concentration was recorded at the mid point of this time interval, except for the initial value, which was taken in the 30 minutes prior to lights on, and the final sample, which was taken in the 30 minutes prior to lights off. Absorbances were read on a Bausch and Lomb Spectronic 20, and calculations of the HCHO concentration from the absorbance and volume of air samples (HCHO vol) were made from the following equation:

$$\text{HCHO (ppm)} = \frac{\text{HCHO } (\mu\text{g}) \times 2.037}{\text{HCHO (vol)}}$$

where HCHO (μg) is taken from the least squares fit of the experimentally determined calibration function of HCHO (μg) vs. absorbance.

The SAMPLE TEMPERATURE was read from either a Doric Thermocouple indicator (°F), using a thermocouple suspended in the chamber (TS2), or from a 19-35°C (0.01 degree/division) thermometer hung free inside the chamber close to the end window, but not in the direct light path (TS1).

RELATIVE HUMIDITY (RH) was measured using a Brady array (Thunder Scientific). The response in volts (V) was converted to percent RH, using

the calibration function supplied by the manufacturer.

HYDROCARBONS (HC) were monitored by gas chromatography with flame ionization detection (GC-FID), using the columns and methods developed by Stephens and Burleson.^{16,17} Methane and C₂ HC's were analyzed using a 5' Poropak N Column, C₃-C₆ HC's using a 36' 2,4-dimethyl sulfolane column, and aromatics and oxygenates using a special three-part column. Oxygenates were also monitored using a 10' Carbowax 600 column. Each GC was calibrated frequently using specially prepared samples.¹⁶ Computer processing of the data includes calculation of the concentration in ppbC for each data point.

B. Experimental Procedures

General. Following each experiment in this program, the glass chamber was flushed with dry air provided by an air purification system¹⁸ (see Figure 2) for about 2 hours at a flow of ~12 cfm. The chamber was then flushed with humidified pure air for about one hour just prior to the start of a run to achieve the desired initial RH. The temperature of the chamber prior to turning on the lamps was adjusted to the operating temperature anticipated during the irradiation by means of infrared lamps. During all flushing procedures, the two sonic pumps were in operation to provide maximum release of materials from the chamber walls.

The matrix air used during the flushing procedure and for the final fill for the experiment generally contained less than a total 60 ppbC of all hydrocarbons except methane, which was typically at a concentration between 550-850 ppb.¹⁸ After completion of filling, analysis of the matrix air prior to injections showed somewhat higher hydrocarbon values due to off-gassing from the chamber walls, but generally these values were less than 200 ppbC nonmethane hydrocarbon.

Following flushing, starting materials were injected by using 100-ml precision bore syringes or micropipettes, and rapid mixing was obtained by brief (~5 minutes) use of the sonic pumps. During the run, the sample temperature was controlled at 32 ± 2°C by means of a variable air flow past the chamber walls.

Light intensity (k₁) was periodically determined, using the method of Holmes et al.,¹⁹ which employs the initial rate of NO₂ photolysis in




Figure 2. Air Purification System.

N_2 as a measure of absolute photon flux in the glass chamber in the actinic region (300-450 nm). However, some O_2 (of order tens of ppm or higher) will be present in the chamber which cannot be evacuated, but instead must be flushed repeatedly with N_2 . Oxygen present in concentrations greater than approximately 10 ppm will lead to somewhat low values for k_1 as they were calculated in this study relative to the true k_1 which would be observed for the complete absence of O_2 . Based on the purity of the nitrogen used (with respect to oxygen content) and a calculated efficiency for repetitive flushing of the chamber, it is estimated that the observed k_1 could be as much as 25% low with respect to the true k_1 .

At full light intensity a k_1 of 0.41 min^{-1} was obtained and at 70% of maximum intensity, 0.26 min^{-1} .

Half-lives for ozone decay, measured at various times in this program under standard conditions of temperature, relative humidity, and light intensity, were ≥ 25 hours in the dark and 12 to 15 hours with irradiation. These half-lives are significantly longer than those reported for other chambers with comparable dimensions, surface to volume ratios, and light intensities and result in correspondingly smaller perturbations on ozone rates of formation and maximum ozone concentrations in the glass chamber experiments.

OH Rate Determinations. A variety of special surrogate mixtures were used in which the liquid components of the original surrogate (2-methyl-2-butene, 2,3-dimethylbutane, toluene and m-xylene) were replaced with the desired quantities of the new compounds to be studied. Isobutene was also added in some cases for use as a reference compound. A gaseous surrogate mixture containing no n-butane was used in the determination of rate of disappearance of methyl ethyl ketone. Hydrocarbons were injected into the chamber from previously prepared mixtures: the liquid hydrocarbons were prepared as a gaseous mixture in a nitrogen-filled 20-l Pyrex bottle or as a liquid mixture, and the gaseous hydrocarbons were contained in an LPO bottle pressurized with nitrogen.

Experiments were carried out at either 70% or 100% of the maximum possible light intensity. Corrections for losses due to sampling were made based on the measured sampling rates of all instruments.

Sulfur Dioxide Decay Experiments. Temperature control for the

experiments carried out without irradiation was accomplished by incremental adjustments of six infrared lamps positioned on three sides of the chamber. During the day the temperature was generally maintained at $33 \pm 1^\circ\text{C}$ and at $33 \pm 2^\circ\text{C}$ overnight. For experiments with irradiation, the temperature was maintained by adjustment of the rate of flow of cooled air between the lamp banks and the chamber walls. The amount of cooling of the air was also varied to provide slight temperature adjustment. However, for experiments carried out at the full irradiation intensity, the lowest chamber temperature which can be maintained with maximum cooling and air flow is about 33°C . At these temperatures the maximum relative humidity which can be employed without condensation in the sample lines and/or instruments (which are at room temperature) is about 50% RH.

To accurately determine very low SO_2 rates during chamber experiments, a precise knowledge of the dilution (due to sampling and chamber leaks) rate is necessary. In an attempt to obtain this information, difluorodichloromethane (FC-12), a chemically inert compound, was added and its disappearance was monitored accurately by gas chromatography (gc). This method is limited, however, by the discovery that isobutane has the same gc retention time under the conditions used as does FC-12. Since isobutane is one of the major ambient hydrocarbons, on days of moderate air pollution the loss of FC-12 could be counteracted by diffusion of ambient isobutane into the chamber. Thus, the observed dilution rates may be low due to interference by isobutane.

III. DEVELOPMENT OF A REACTIVITY SCALE AND CLASSIFICATION FOR ORGANICS BASED ON THEIR RATES OF REACTION WITH THE HYDROXYL RADICAL

A. Relative Rates of Reaction of Organic Compounds with the Hydroxyl Radical

The use of the relative rates of reaction of hydrocarbons determined under simulated atmospheric conditions using the SAPRC all-glass environmental chamber has been previously validated as a method of determining rate constants for reaction of hydrocarbons with the hydroxyl radical (OH).^{20,21} We have now extended the application of this chamber method to the determination of OH rate constants for more than 30 additional compounds.²²⁻²⁶ The assumption that reaction with the OH radical is by far the dominant reaction pathway for consumption of hydrocarbons during the first 2-3 hours of irradiation in these experiments (conducted at 1 atm pressure in air) has been confirmed by the good agreement observed between OH rate constants measured by the chamber method and those determined in elementary rate studies of the individual hydrocarbon. This agreement between relative and absolute rate determinations is illustrated in Figure 3. The extent to which this assumption is valid is also supported by the results of computer modeling calculations²⁷ (shown in Figure 4) for an HC-NO_x system of total hydrocarbon concentration identical to that used in one of these studies.²² In the computer simulation, a propene and n-butane mixture was used as a surrogate for the complex hydrocarbon mixture employed in the chamber experiment, and the rate of attack on propene by OH, O₃, O(³P), and HO₂ was calculated. The relative and total concentrations of propene and n-butane were chosen such that the overall hydrocarbon reactivity towards the OH radical would equal that predicted for the complex mixture. It is clear from Figure 4 that although OH is the major attacking species in these experiments, the O₃ contribution to the disappearance rates of the alkenes increases with the time of irradiation. In contrast, the rates of reaction of O₃ with alkanes and aromatics are many orders of magnitude slower,²⁸⁻³² and no correction for their reaction with ozone was necessary. During the initial hours of irradiation other species such as NO₃³³ and HO₂³⁴ may contribute slightly to hydrocarbon disappearance rates, especially for alkenes, but since




Figure 3. Comparison of Relative Rates of Hydrocarbon Disappearance Determined by the Chamber Method with Selected Published Rate Constants for Reaction of Those Hydrocarbons with OH Radicals.




Figure 4. Predicted Relative Importance of Several Reactive Intermediates during Photooxidation of a Propene-n-Butane Mixture under Simulated Atmospheric Conditions.

their concentrations, and in some cases rate constants, are not known, correction is not possible.

Results. As had been proposed, four general types of compounds, based on their source or potential atmospheric significance, were studied: combustion source hydrocarbons, solvent components, naturally occurring hydrocarbons, and several compounds which are believed to be potential organic aerosol precursors (see Table 5). The relative rates of disappearance of some selected compounds are shown in Figures 5-7. Absolute rate constants were derived from the relative rates of disappearance using the literature values for the absolute OH rate constants for either n-butane³⁵⁻⁴¹ or isobutene⁴² (1.8×10^9 and 3.05×10^{10} l mole⁻¹ sec⁻¹, respectively). The compounds by hydrocarbon class and the rate constants thus derived are given in Table 6.

Detailed discussions of the significance, both fundamental and applied, of these rate constant determinations as well as calculations of atmospheric half-lives and analyses of the role of these compounds in atmospheric chemistry have already been reported in the literature²²⁻²⁶ and reprints are included in Appendix A.

B. Development of a Reactivity Scale and Reactivity Classification Based on the Reaction of Organics with OH

The important role of OH in photochemical smog formation is now well established⁴³⁻⁵⁰ on the basis of both experimental and computer modeling studies, and OH has been directly observed⁵¹⁻⁵³ in both the laboratory and in ambient air. This awareness has led to a dramatic increase during the past several years in the number of compounds for which OH rate constants have been measured. In addition to the extensive elementary rate determinations carried out in this laboratory^{35,54-62} and elsewhere, we have, as described above and previously,²¹ measured rate constants for 40 compounds using the chamber method. This technique has also been employed recently by Niki and coworkers.⁶³

From the successful correlation of OH rate constants with the rates of hydrocarbon disappearance observed in chamber simulations at the SAPRC^{20,22-26,62} and Ford Motor Co.⁶³ laboratories, we conclude that, to a good approximation, this correlation can be extrapolated to the atmosphere

Table 5. Rate Constants for Reaction of OH with Organic Compounds
Obtained Using the Chamber Method

Compound	Rate Constant 10^{-9} (ℓ mole $^{-1}$ sec $^{-1}$)	Reference
Combustion Sources		
Ethene	5.2 ± 1.0	a
Propene	17.5 ± 3.5	a
cis-2-Butene	14.9 ± 2.2	b
	39.2 ± 8.0	a
1,3-Butadiene	37.9 ± 5.6	b
	32.9 ± 6.6	c
2,3-Dimethylbutane	46.4 ± 9.3	a
Isopentane	3.1 ± 0.5	f
2-Methylpentane	2.0 ± 0.4	a
3-Methylpentane	3.2 ± 0.6	a
n-Hexane	4.3 ± 0.9	a
m-Xylene	3.8 ± 0.8	a
n-Propylbenzene	12.9 ± 2.6	a
Isopropylbenzene	3.7 ± 0.8	a
Ethylbenzene	3.7 ± 0.8	a
o-Ethyltoluene	4.8 ± 1.0	a
m-Ethyltoluene	8.2 ± 1.6	a
p-Ethyltoluene	11.7 ± 2.3	a
	7.8 ± 1.6	a
Solvent Components		
Methyl ethyl ketone	2.0 ± 0.6	b
Methyl isobutyl ketone	9.0 ± 3.0	b
Diisobutyl ketone	15 ± 5	b
Isopropyl alcohol	4.3 ± 1.3	d
Diethyl ether	4.3 ± 1.3	d
Di-n-propyl ether	5.6 ± 1.1	d
Tetrahydrofuran	10.4 ± 2.1	d
n-Propyl acetate	8.8 ± 1.8	e
sec-Butyl acetate	2.6 ± 0.5	e
	3.4 ± 0.7	e
Naturally Occurring Hydrocarbons		
α -Pinene	34.8 ± 5.2	b
β -Pinene	40.6 ± 6.1	b
d-Limonene	90.0 ± 13.5	b
Potential Aerosol Precursors		
2,2,3-Trimethylbutane	2.3 ± 0.5	f
1-Heptene	22 ± 5	f
Cyclohexene	47 ± 9	f
1-Methylcyclohexene	58 ± 12	f

^aLloyd, Darnall, Winer and Pitts, J. Phys. Chem., 80, 789 (1976).

^bWiner, Lloyd, Darnall and Pitts, J. Phys. Chem., 80, 1635 (1976).

^cDarnall, Winer, Lloyd and Pitts, unpublished results, 1976.

^dLloyd, Darnall, Winer and Pitts, Chem. Phys. Lett., 42, 205 (1976)

^eWiner, Lloyd, Darnall and Pitts, Chem. Phys. Lett., submitted (1977).

^fDarnall, Winer, Lloyd and Pitts, Chem. Phys. Lett., 44, 415 (1976).




Figure 5. Concentrations of Selected Organic Compounds during Irradiation of a HC-NO_x mixture in Air at 305 ± 2 K (Propene and 1,3-Butadiene Concentrations Have Been Corrected for Reaction with Ozone).




Figure 6. Concentrations of Ketones and Isobutene during Irradiation of a HC-NO_x Mixture in Air at 305 ± 2 K.




Figure 7. Concentrations of Ethers and Isobutene during Irradiation of a HC-NO_x Mixture in Air at 305 ± 2 K.

Table 6. Rate Constants for Reaction of OH with Organic Compounds Obtained Using the Chamber Method

Compound	Rate Constant 10^{-9} (ℓ mole $^{-1}$ sec $^{-1}$)	Reference
<u>Alkenes</u>		
Ethene	5.2 \pm 1.0	a
Propene	17.5 \pm 3.5	a
cis-2-Butene	14.9 \pm 2.2	b
	39.2 \pm 8.0	a
	37.9 \pm 5.6	b
	32.9 \pm 6.6	c
1,3-Butadiene	46.4 \pm 9.3	a
1-Heptene	22 \pm 5	d
Cyclohexene	47 \pm 9	d
1-Methylcyclohexene	58 \pm 12	d
α -Pinene	34.8 \pm 5.2	b
β -Pinene	40.6 \pm 6.1	b
d-Limonene	90.0 \pm 13.5	b
<u>Alkanes</u>		
2,3-Dimethylbutane	3.1 \pm 0.5	d
Isopentane	2.0 \pm 0.4	a
2-Methylpentane	3.2 \pm 0.6	a
3-Methylpentane	4.3 \pm 0.9	a
n-Hexane	3.8 \pm 0.8	a
2,2,3-Trimethylbutane	2.3 \pm 0.5	d
<u>Aromatic Hydrocarbons</u>		
m-Hylene	12.9 \pm 2.6	a
n-Propylbenzene	3.7 \pm 0.8	a
Isopropylbenzene	3.7 \pm 0.8	a
Ethylbenzene	4.8 \pm 1.0	a
o-Ethyltoluene	8.2 \pm 1.6	a
m-Ethyltoluene	11.7 \pm 2.3	a
p-Ethyltoluene	7.8 \pm 1.6	a
<u>Oxygenated Compounds</u>		
Methyl ethyl detone	2.0 \pm 0.6	b
Methyl isobutyl ketone	9.0 \pm 3.0	b
Diisobutyl ketone	15 \pm 5	b
Isopropyl alcohol	4.3 \pm 1.3	e
Diethyl ether	5.6 \pm 1.1	e
Di-n-propyl ether	10.4 \pm 2.1	e
Tetrahydrofuran	8.8 \pm 1.8	f
n-Propyl acetate	2.6 \pm 0.5	f
sec-Butyl acetate	3.4 \pm 0.7	f

^aLloyd, Darnall, Winer and Pitts, J. Phys. Chem., 80, 789 (1976).

^bWiner, Lloyd, Darnall and Pitts, J. Phys. Chem., 80, 1635 (1976).

^cDarnall, Winer, Lloyd and Pitts, unpublished results, 1976.

^dLloyd, Darnall, Winer and Pitts, Chem. Phys. Lett., 42, 205 (1976).

^eWiner, Lloyd, Darnall and Pitts, Chem. Phys. Lett., submitted (1977).

^fDarnall, Winer, Lloyd and Pitts, Chem. Phys. Lett., 44, 415 (1976).

(a) for alkenes in ambient air parcels during the early morning hours when ozone levels are generally quite low (≤ 0.05 ppm), and (b) for alkanes, aromatics, and most of the other classes of compounds discussed here at essentially all times and locations. The latter assumption, namely that an OH rate constant is a good "reactivity index" for alkanes, aromatics, halocarbons, etc. throughout an irradiation day (or multiple irradiation days) rests upon the fact that the rates of reaction of these classes of organics with species such as ozone, $O(^3P)$ atoms and the hydroperoxyl radical HO_2 are substantially slower than with OH.^{28-30,64,65} For example, even at the highest ozone concentrations experienced in ambient atmospheres, O_3 will not contribute significantly to the consumption of alkanes and aromatics. This is in contrast to the case for alkenes which, although the rate constants for reaction of O_3 are not particularly large,^{31,32,65} are depleted rapidly by reaction with ozone at the average concentrations commonly encountered in polluted ambient air (~ 0.1 to 0.2 ppm).

Proposed Reactivity Scale and Classification. In view of the above, and the very recent availability of a large number of OH rate constants, we have proposed^{66,67} a hydrocarbon reactivity scale based on the primary chemical act of hydroxyl radical (OH) attack on organic species. We have done so on two bases. First, as discussed above, the OH radical is a key reactive intermediate in photochemical air pollution. Second, a reactivity scale based on the depletion of hydrocarbons by reactions with OH has utility in assessing hydrocarbon chemical behavior in polluted ambient air, since only those compounds which participate at "significant" rates in atmospheric reactions are of consequence in the chemical transformations in ambient air, including, of course, the production of photochemical oxidant. Thus, our working premise has been that the relative reactivity of organics towards OH is generally a useful and directly measurable index of their potential importance in the production of secondary pollutants. As we have previously pointed out,⁶⁸ our proposed reactivity classification would not be valid if specific reactivity criteria unrelated in varying degrees to oxidant formation (for example, aerosol formation and associated visibility effects, mutagenic activity, etc.) are of concern.

Although a reactivity scale based on OH rate constants can, in principle, be divided into a large number of classes (see below), it is

convenient, particularly for purposes of comparison with other scales, to use a five-class scale which is based on order of magnitude differences in reactivities of organics toward OH relative to methane (=1). This scale is shown in Table 7, which also gives the range of half-lives of compounds appearing in each class assuming depletion solely due to reaction with the OH radical at a concentration of 10^7 radicals cm^{-3} .

Table 7. Reactivity Scale for Organics Based on Rate of Consumption Due to Reaction with the Hydroxyl Radical

Class	Half-Life ^a (days)	Reactivity Relative to Methane (=1)
I	≥ 10	≤ 10
II	1 - 10	10 - 100
III	0.1 - 1	100 - 1000
IV	0.01 - 0.1	1000 - 10,000
V	≤ 0.01	$\geq 10,000$

^a $t_{1/2} = 0.0693/k_{\text{OH}}[\text{OH}]$. $[\text{OH}]$ is assumed to be 10^7 radicals cm^{-3} .

We have attempted to present a reactivity classification which is exhaustive with respect to the reported room temperature literature for OH rate constant determinations (as well as recent, as yet unpublished, measurements in this laboratory). Thus, we have compiled a reactivity classification consisting of more than 110 compounds assigned into the 5 classes of our proposed reactivity scale.⁶⁹ Table 8 gives the rate constant (with its source)*, the reactivity (relative to methane = 1)

* The rate constant data given in Table 8 are not to be construed as critically evaluated or recommended values. In general, absolute rate determinations were used when available, and average values are often given.

Table 8. Rate Constants for Reaction of Organics (and Selected Inorganics) with the Hydroxyl Radical and the Corresponding Reactivity Classification

Compound	$k_{OH}^a + \text{Cpd}$ ($\ell \text{ mol}^{-1} \text{ s}^{-1}$) $\times 10^{-9}$	Reference	Reactivity Relative to Methane = 1	Proposed Class
<u>Alkanes</u>				
Methane	0.0048	b-f	1	I
Ethane	0.17	c, f, g	35	II
Propane	1.3	f, h	270	III
n-Butane	1.8	c, h-m	375	III
Isobutane	1.3	c, n	270	III
Cyclobutane	0.72	h	150	III
n-Pentane	3.9	n	810	III
Isopentane	2.0	o	420	III
Cyclopentane	3.7	p	765	III
Neopentane	0.49	c	100	III
n-Hexane	3.6	n, o, q	750	III
Cyclohexane	4.2	c, h, n	875	III
2-Methylpentane	3.2	o	665	III
3-Methylpentane	4.3	o	900	III
2,3-Dimethylbutane	3.1	r	645	III
2,2,3-Trimethylbutane	2.3	r	480	III
2,2,4-Trimethylpentane	2.3	c	480	III
n-Octane	5.1	c	1060	IV
2,2,3,3-Tetramethylbutane	0.67	c	140	III
<u>Alkenes</u>				
Ethene	4.7	s	985	III
Propene	15.1	t	3145	IV
1-Butene	21.3	t	4440	IV
Isobutene	30.5	t	6355	IV
cis-2-Butene	32.3	t	6730	IV
trans-2-Butene	42.1	t	8770	IV
1-Pentene	18	n	3750	IV
cis-2-Pentene	39	n	8125	IV
2-Methyl-1-butene	35	n	7290	IV
2-Methyl-2-butene	48	u	10,000	V
1-Hexene	19	n	3960	IV
Cyclohexene	43	r, n	8960	IV
2,3-Dimethyl-2-butene	92	l	19,170	V
3,3-Dimethyl-1-butene	17	n	3540	IV
1-Methylcyclohexene	58	r	12,080	V
1-Heptene	22	r	4585	IV
<u>Di-Alkenes</u>				
Allene	2.7	v	560	III
1,3-Butadiene	46.4	o	9670	IV
Isoprene	47	w	9790	IV

(continued)

Table 8. (Continued)

Compound	$k_{OH + Cpd}^a$ ($\ell \text{ mol}^{-1} \text{ s}^{-1}$) $\times 10^{-9}$	Reference	Reactivity Relative to Methane = 1	Proposed Class
<u>Alkynes</u>				
Acetylene	0.099	x	21	II
Methylacetylene	0.57	v	120	III
<u>Aromatic Hydrocarbons</u>				
Benzene	0.85	y,z	180	III
Toluene	3.6	y,z	750	III
n-Propylbenzene	3.7	o	770	III
Isopropylbenzene	3.7	o	770	III
Ethylbenzene	4.8	o	1000	IV
p-Xylene	7.4	y,aa	1540	IV
o-Xylene	8.4	y,aa	1750	IV
m-Xylene	14.1	y,aa	2940	IV
p-Ethyltoluene	7.8	o	1625	IV
o-Ethyltoluene	8.2	o	1710	IV
m-Ethyltoluene	11.7	o	2440	IV
p-Cymene	9.2	w	1920	IV
1,2,3-Trimethylbenzene	14.9	y,aa	3100	IV
1,2,4-Trimethylbenzene	20	y,aa	4170	IV
1,3,5-Trimethylbenzene	29.7	y,aa	6190	IV
<u>Ketones</u>				
Methyl ethyl ketone	2.0	w	420	III
Methyl isobutyl ketone	9.0	w	1875	IV
Diisobutyl ketone	15	w	3125	IV
<u>Alcohols</u>				
Methanol	0.57	q	120	III
Ethanol	1.8	q	375	III
n-Propyl alcohol	2.3	q	480	III
Isopropyl alcohol	4.3	bb	895	III
n-Butanol	4.1	q	855	III
4-Methyl-2-pentanol	4.3	cc	895	III
<u>Ethers</u>				
Diethyl ether	5.6	bb	1170	IV
Di-n-propyl ether	10.4	bb	2170	IV
Tetrahydrofuran	8.8	dd	1830	IV
<u>Esters</u>				
n-Propyl acetate	2.7	dd	560	III
sec-Butyl acetate	3.4	dd	710	III

(continued)

Table 8. (Continued)

Compound	$k_{OH + Cpd}^a$ ($\ell \text{ mol}^{-1} \text{ s}^{-1}$) $\times 10^{-9}$	Reference	Reactivity Relative to Methane = 1	Proposed Class
<u>Natural Hydrocarbons</u>				
α -Pinene	35	w	7290	IV
β -Pinene	41	w	8540	IV
p-Menthane	4	w	830	III
3-Carene	52	w	10,830	V
β -Phellandrene	70	w	14,580	V
Carvomenthene	76	w	15,830	V
d-Limonene	90	w	18,750	V
Dihydromyrcene	101	w	21,040	V
Myrcene	137	w	28,540	V
cis-Ocimene	192	w	40,000	V
<u>Halocarbons</u>				
Fluoromethane	0.0096	d	2	I
Chloromethane	0.025	d, ee, ff	5	I
Bromomethane	0.023	d, ff	5	I
Difluoromethane	0.0047	d	1	I
Chlorofluoromethane	0.022	d	5	I
Dichloromethane	0.084	d, ee, ff	18	II
Trifluoromethane	0.00012	d	0.025	I
Chlorodifluoromethane	0.025	d, gg	5	I
Dichlorofluoromethane	0.016	d, ee	3	I
Trichloromethane	0.065	d, ff	14	II
Tetrafluoromethane	<0.00024	d	<0.05	I
Chlorotrifluoromethane	<0.00042	d	<0.09	I
Dichlorodifluoromethane	<0.0006	d, gg	<0.13	I
Fluorotrifluoromethane	<0.0006	d, gg	<0.13	I
Tetrachloromethane	<0.0024	d	<0.5	I
Chloroethane	0.235	g	49	II
1,2-Dichloroethane	0.132	g	28	II
1,2-Dibromoethane	0.15	g	31	II
1,1-Difluoroethane	0.019	g	4	I
1,1-Dichloroethane	0.16	g	33	II
1-Chloro-1,1-difluoroethane	0.0017	g	0.35	I
1,1,1-Trichloroethane	0.0090	g	2	I
1,1,1-Trifluoro-2-chloroethane	0.0063	g	1	I
1,1,1,2-Tetrafluoro-2-chloroethane	0.0075	g	2	I
1,1,1-Trifluoro-2,2-dichloroethane	0.0171	g	4	I
1,1,2,2-Tetrafluoro-1,2-dichloroethane	<0.0003	g	<0.06	I
1,1,2-Trifluoro-1,2,2-trichloroethane	<0.0002	g	<0.04	I

(continued)

Table 8. (Continued)

Compound	$k_{\text{OH} + \text{Cpd}}^{\text{a}}$ ($\ell \text{ mol}^{-1} \text{ s}^{-1}$) $\times 10^{-9}$	Reference	Reactivity Relative to Methane = 1	Proposed Class
<u>Nitrogen-Containing Compounds</u>				
Ammonia	0.0988	hh	22	II
Methylamine	13.2	ii	2750	IV
<u>Sulfur-Containing Compounds</u>				
Hydrogen sulfide	3.16	hh	660	III
Methyl mercaptan	20.4	ii	4250	IV
<u>Miscellaneous</u>				
Carbon monoxide	0.084	jj	18	II
Methoxybenzene	11.8	ii	2460	IV
o-Cresol	20.5	ii	4280	IV

^a "At room temperature" (i.e., within range 292-305 K).

^b D. D. Davis, S. Fischer, and R. Schiff, J. Chem. Phys., 61, 2213 (1974).

^c N. R. Greiner, J. Chem. Phys., 53, 1070 (1970).

^d C. J. Howard and K. M. Evenson, J. Chem. Phys., 64, 197 (1976).

^e J. J. Margitan, F. Kaufman, and J. G. Anderson, Geophys. Res. Lett., 1, 80 (1974).

^f R. P. Overend, G. Paraskevopoulos, and R. J. Cvetanovic, Can. J. Chem., 53, 3374 (1975).

^g C. J. Howard and K. M. Evenson, J. Chem. Phys., 64, 4303 (1976).

^h R. A. Gorse and D. H. Volman, J. Photochem., 3, 115 (1974).

ⁱ R. A. Perry, R. Atkinson and J. N. Pitts, Jr., J. Chem. Phys., 64, 5314 (1976).

^j I. M. Campbell, B. J. Handy, and R. M. Kirby, J. Chem. Soc. Faraday Trans., I 71, 867 (1975).

^k S. Gordon and W. A. Mulac, Intern. J. Chem. Kinetics, Symposium No. 1, 289 (1975).

^l E. D. Morris, Jr. and H. Niki, J. Phys. Chem., 75, 3640 (1971).

^m F. Stuhl, Z. Naturforsch. 28A, 1383 (1973).

ⁿ C. H. Wu, S. M. Japar, and H. Niki, J. Environ. Sci. Health, A11, 191 (1976).

^o A. C. Lloyd, K. R. Darnall, A. M. Winer, and J. N. Pitts, Jr., J. Phys. Chem., 80, 789 (1976).

(continued)

Table 8. (Continued)

- ^p D. H. Volman, Intern. J. Chem. Kinetics, Symposium No. 1, 358 (1975).
- ^q I. M. Campbell, D. R. McLaughlin, and B. J. Handy, Chem. Phys. Lett., 38, 362 (1976).
- ^r K. R. Darnall, A. M. Winer, A. C. Lloyd, and J. N. Pitts, Jr., Chem. Phys. Lett., submitted for publication (1976).
- ^s R. Atkinson, R. A. Perry, and J. N. Pitts, Jr., J. Chem. Phys., submitted for publication (1976).
- ^t R. Atkinson and J. N. Pitts, Jr., J. Chem. Phys., 63, 3591 (1975).
- ^u R. Atkinson, R. A. Perry, and J. N. Pitts, Jr., Chem. Phys. Lett., 38, 607 (1976).
- ^v J. N. Bradley, W. Hack, K. Hoyermann, and H. Gg. Wagner, J. Chem. Soc. Faraday Trans. I 69, 1889 (1973).
- ^w A. M. Winer, A. C. Lloyd, K. R. Darnall, and J. N. Pitts, Jr., J. Phys. Chem., 80, 1635 (1976).
- ^x D. D. Davis, S. Fischer, R. Schiff, R. T. Watson, and W. Bollinger, J. Chem. Phys., 63, 1707 (1975).
- ^y D. A. Hansen, R. Atkinson, and J. N. Pitts, Jr., J. Phys. Chem., 79, 1763 (1975).
- ^z D. D. Davis, W. Bollinger, and S. Fischer, J. Phys. Chem., 79, 293 (1975).
- ^{aa} G. J. Doyle, A. C. Lloyd, K. R. Darnall, A. M. Winer, and J. N. Pitts, Jr., Environ. Sci. Technol., 9, 237 (1975).
- ^{bb} A. C. Lloyd, K. R. Darnall, A. M. Winer, and J. N. Pitts, Jr., Chem. Phys. Lett., 42, 205 (1976).
- ^{cc} J. L. Laity, I. G. Burstein, and B. R. Appel, Advan. Chem. Series, 124, 95 (1973).
- ^{dd} A. M. Winer, A. C. Lloyd, K. R. Darnall, and J. N. Pitts, Jr., in preparation.
- ^{ee} R. A. Perry, R. Atkinson, and J. N. Pitts, Jr., J. Chem. Phys., 64, 1618 (1976).
- ^{ff} D. D. Davis, G. Machado, B. Conaway, Y. Oh, and R. Watson, J. Chem. Phys., 65, 1268 (1976).
- ^{gg} R. Atkinson, D. A. Hansen, and J. N. Pitts, Jr., J. Chem. Phys., 63, 1703 (1975).
- ^{hh} R. A. Perry, R. Atkinson, and J. N. Pitts, Jr., J. Chem. Phys., 64, 3237 (1976).
- ⁱⁱ R. A. Perry, R. Atkinson, and J. N. Pitts, Jr., unpublished results, (1976).
- ^{jj} R. F. Hampson, Jr. and D. Garvin, Eds., NBS Technical Note 866, June, 1975.

and proposed classification for each compound, with the compounds arranged by chemical class. In view of the growing attention being given to nitrogen- and sulfur-containing compounds in the polluted troposphere we have included ammonia, methylamine, hydrogen sulfide, and methyl mercaptan in our classification (as well as carbon monoxide as a matter of interest). For convenience, we also present, in Table 9, the compounds appearing in each of the 5 classes of the SAPRC reactivity scale in order of increasing reactivity within each class.

Comparison with Other Reactivity Classification. In our proposed reactivity classification Class I contains methane and most halocarbons. The remaining half-dozen halocarbons appear in Class II along with carbon monoxide, ammonia, acetylene, and ethane. Almost all alkanes appear in Class III, while alkenes other than ethene appear in Classes IV or V. Aromatics and oxygenates appear in Classes III or IV and the natural hydrocarbons (not specifically identified in previous reactivity classifications) are of high reactivity and appear in Classes IV and V. Clearly our proposed classification emphasizes that most compounds react in polluted atmospheres and Classes I and II contain the relatively few compounds which have half-lives greater than 10 days and 24 hours, respectively. This result parallels the conclusion by Altshuller and Bufalini in their 1971 review article that "almost every hydrocarbon except methane can produce some oxidant when photooxidized in the presence of high enough ratios of hydrocarbons to oxides of nitrogen."

Although our proposed reactivity scale and classification is based solely on consumption rates of organics, Altshuller and Bufalini⁷⁰ have shown that this measure of reactivity is similar to one based on nitric oxide oxidation rates. They showed that the ranking of reactivities of hydrocarbons from the nitric oxide photooxidation studies of Altshuller and Cohen⁷¹ and Glasson and Tuesday⁶⁴ was essentially the same as that obtained from the studies of hydrocarbon consumption carried out by Schuck and Doyle,⁷² Stephens and Scott,⁷³ and Tuesday.⁷⁴ The ranking of reactivities for the aromatic hydrocarbons in our classification is very similar to that obtained by Altshuller et al.⁷⁵ and by Kopczynski.^{76,77} We have previously^{66,67} shown that our classification is in general agreement with that proposed by Dimitriades in 1974⁷⁸ and noted where significant

Table 9. Proposed Reactivity Classification of Organics (and Selected Inorganics) Based on Reaction with the Hydroxyl Radical

Class I ($\leq 4.8 \times 10^7$) ^a	Class II ($4.8 \times 10^7 - 4.8 \times 10^8$) ^a	Class III ($4.8 \times 10^8 - 4.8 \times 10^9$) ^a	Class IV ($4.8 \times 10^9 - 4.8 \times 10^{10}$) ^a	Class V ($\geq 4.8 \times 10^{10}$) ^a
Trifluoromethane	Trichloromethane	Neopentane	Ethylbenzene	2-Methyl-2-butene
1,1,2-Trichloro-1,2,2-trifluoroethane	Dichloromethane	Methanol	n-Octane	3-Carene
Tetrafluoromethane	Carbon monoxide	Methylacetylene	Diethyl ether	1-Methylcyclohexene
1,2-Dichloro-1,1,2,2-tetrafluoroethane	Ammonia	2,2,3,3-Tetramethylbutane	p-Xylene	β -Phellandrene
Chlorotrifluoromethane	Acetylene	Cyclobutane	p-Ethyltoluene	Carvomenthene
Dichlorodifluoromethane	1,2-Dichloroethane	Benzene	o-Ethyltoluene	d-Limonene
Fluorotrichloromethane	1,2-Dibromoethane	Isobutane	o-Xylene	2,3-Dimethyl-2-butene
1-Chloro-1,1-difluoroethane	Ethane	Propane	Tetrahydrofuran	Dihydromyrcene
Tetrachloromethane	1,1-Dichloroethane	n-Butane	Methyl isobutyl ketone	Myrcene
Difluoromethane	Chloroethane	Ethanol	p-Cymene	cis-Occimene
Methane		Methyl ethyl ketone	Di-n-propyl ether	
2-Chloro-1,1,1-trifluoroethane		Isopentane	m-Ethyltoluene	
2-Chloro-1,1,1,2-tetrafluoroethane		1-Propanol	Methoxybenzene	
1,1,1-Trichloroethane		2,2,4-Trimethylpentane	Methylamine	
Fluoromethane		2,2,3-Trimethylbutane	n-Xylene	
Dichlorofluoromethane		n-Propyl acetate	1,2,3-Trimethylbenzene	
2,2-Dichloro-1,1,1-trifluoroethane		Allene	Diisobutyl ketone	
1,1-Difluoroethane		2,3-Dimethylbutane	Propene	
Chlorofluoromethane		Hydrogen sulfide	3,3-Dimethyl-1-butene	
Bromomethane		2-Methylpentane	1-Pentene	
Chloromethane		sec-Butyl acetate	1-Hexene	
Chlorodifluoromethane		Toluene	1,2,4-Trimethylbenzene	
		Cyclopentane	Methyl mercaptan	
		n-Propylbenzene	o-Cresol	
		Isopropylbenzene	1-Butene	
		n-Hexane	1-Heptene	
		n-Pentane	1,3,5-Trimethylbenzene	
		p-Menthane	Isobutene	
		1-Butanol	cis-2-Butene	
		Cyclohexane	2-Methyl-1-butene	
		2-Propanol	α -Pinene	
		4-Methyl-2-pentanol	cis-2-Pentene	
		3-Methylpentane	β -Pinene	
		Ethene	trans-2-Butene	
			Cyclohexene	
			1,3-Butadiene	
			Isoprene	

^aRange of values (in liter mole⁻¹ sec⁻¹) for the rate constant for reaction of the OH radical with the listed compounds.

differences existed. Several of these differences arose from the greater degree of differentiation in reactivity permitted by the OH reactivity criterion.

A three-class reactivity classification of organic compounds very recently adopted by the ARB⁷⁹ is shown in Table 10, and is similar to one now under consideration by the EPA.⁸⁰ As described by the ARB,^{79,80} Class I includes low reactivity organic compounds, yielding little if any ozone under urban conditions. Class II consists of moderately reactive organic compounds which give an intermediate yield of ozone within the first day of solar irradiation, while Class III is limited to highly reactive organic compounds which give high yields of ozone within a few hours of irradiation.

In general, the ARB three-tiered classification is consistent with the reactivity classification presented here, based on OH rate constants, with only minor exceptions. For example, the ARB scale shows the primary and secondary C₂₊ alcohols to be highly reactive in Class III, while our scale shows them to be of moderate reactivity.

Advantages and Limitations of the OH Reactivity Classification. In contrast to previous reactivity scales, where it is commonly recognized that uncertainties in the number of classes into which the reactivity scales can be meaningfully divided,^{78,80} OH reactivity as developed here could be divided into a large number of classes based on the fact that the accuracy of OH rate constant measurements is generally better than ±20%. Thus, one significant strength of this approach is the very high degree of resolution obtained in ranking compounds in order of their rate of reaction with OH. A second advantage of the present classification is that it can be readily extended to include additional organic (and inorganic) compounds once their rate of reaction with OH is known. Thus, it is not necessary to conduct a major new experimental program each time a significant number of new compounds become of interest (as has been felt necessary in the usual smog chamber approach, in order to obtain consistent data for all compounds of interest at a given time). Finally, our proposed classification gives greater weight than most previous classifications to the alkanes and aromatic hydrocarbons, which as discussed above, require a longer irradiation period but can, during downwind transport or stagnant air episodes, contribute significantly to ozone formation.

Table 10. California Air Resources Board (ARB) Reactivity Classification of Organic Compounds[†]

Class I (Low Reactivity)	Class II (Moderate Reactivity)	Class III (High Reactivity)
C ₁ -C ₂ Paraffins	Mono-tert-alkyl-benzenes	All other aromatic hydrocarbons
Acetylene	Cyclic ketones	All olefinic hydrocarbons (including partially halogenated)
Benzene	Alkyl acetates	Aliphatic aldehydes
Benzaldehyde	2-Nitropropane	Branched alkyl ketones
Acetone	C ₃ + Paraffins	Cellosolve acetate
Methanol	Cycloparaffins	Unsaturated ketones
Tert-alkyl alcohols	n-Alkyl ketones	Primary & secondary C ₂ + alcohols
Phenyl acetate	N-Methyl pyrrolidone	Diacetone alcohol
Methyl benzoate	N,N-Dimethyl acetamide	Ethers
Ethyl amines	Alkyl phenols*	Cellosolves
Dimethyl formamide	Methyl phthalates**	Glycols*
Perhalogenated Hydrocarbons		C ₂ + alkyl phthalates**
Partially halogenated paraffins		Other esters**
Phthalic anhydride**		Alcohol amines**
Phthalic acids**		C ₃ + Organic acids + di acid**
Acetonitrile*		C ₃ + di acid anhydrides**
Acetic acid		Formin** (Hexa methylene-tetramine)
Aromatic amines		Terpenic hydrocarbons
Hydroxyl amines		Olefin oxides**
Naphthalene*		
Chlorobenzenes*		
Nitrobenzenes*		
Phenol*		

[†] This reactivity classification is identical to that suggested by the EPA in August, 1975^{79,80} except that the ARB has moved propane from Class I to Class II. *Reactivity data are either non-existent or inconclusive, but conclusive data from similar compounds are available; therefore, rating is uncertain but reasonable. **Reactivity data are uncertain.

These strengths notwithstanding, several caveats should be borne in mind in applying the proposed OH reactivity scale. First of all, it is not strictly applicable to compounds which undergo significant photodissociation in the atmosphere (e.g., aliphatic aldehydes) or which react in the atmosphere at significant rates with species other than OH (e.g., the alkenes with O_3). In such cases, the compound will be more reactive than predicted from a scale based on consumption due solely to OH attack. It is interesting to note, however, that with respect to the reaction of ozone with alkenes, this limitation is not as serious as it would appear to be based only on the relative rates of consumption of alkenes by ozone and OH, respectively. This follows from the fact that recently Niki and coworkers⁸¹ have shown that O_3 -alkene reactions do not lead predominantly to radical products, and hence this additional alkene consumption pathway will contribute significantly less "additional reactivity" (over that obtained for OH reaction alone) than if it led to radical chain propagation.

A second, and perhaps more significant, limitation in the application of our reactivity classification concerns the inherent problem arising from uncertainties in the identity and fates of subsequent products.⁷⁸ For example, two compounds which have the same rate constant for reaction with OH may not necessarily have the same oxidant-producing potential in undergoing further photooxidation. As another example, we have recently reported that alkoxy radicals of the higher alkenes ($\geq C_4$) undergo internal isomerization⁸² and also that reaction of alkylperoxy radicals ($\geq C_4$) with NO leads to the formation of alkyl nitrates ($RONO_2$) rather than to $RO + NO_2$.⁸³ Thus, estimates of ozone formation (or other secondary smog manifestations) for the higher alkanes which are based solely on their rates of reaction with OH are likely to be somewhat high since nitrate formation is effectively radical terminating.

IV. A CHAMBER STUDY OF SULFATE AEROSOL FORMATION

A. Introduction

The formation of inorganic aerosols containing sulfate and nitrate anions is a well established feature of photochemical air pollution. Recent work has reemphasized the overriding importance of these anions for the phenomenon of visibility reduction.⁸⁴⁻⁸⁶ It is also becoming apparent that inhaled sulfate and nitrate aerosols have appreciable effects on the health of exposed populations,^{4,87-89} a more serious although until recently less well-known effect than visibility reduction.

The ultimate fate of emitted sulfur compounds is conversion to sulfate and eventual transfer to the oceans on a worldwide basis. Conversion processes are active throughout the atmosphere⁹⁰ but proceed at an accelerated rate in a polluted urban atmosphere. Accumulation of sulfate is due to two factors, a higher precursor emission rate than the worldwide average, primarily due to the combustion of sulfur-containing fossil fuels, and the promotion of conversion processes by the presence of other anthropogenic primary or secondary pollutants. One major promoted conversion process is the entrainment of anthropogenic sulfur compounds, mostly sulfur dioxide, into the reactions accompanying the nitrogen oxides-promoted photooxidation of organics (mostly hydrocarbons).

In recent years the sulfate burden in the SCAB has been increasing at a rate of about 10% per year.⁵ The inorganic aerosol problem is therefore not under control and promises to become significantly worse in the future, especially if aggravated by nondiscriminatory use of higher sulfur-content fuels as a result of fuel shortages.

The major uncertainties in the above projections are the effects of current and future control programs on the conversion processes. As hydrocarbon emissions are reduced, the atmospheric photochemical activity will presumably decrease, but whether this will reduce conversion rates proportionately remains to be seen. Thus, it is possible that the rates will be reduced only sufficiently to transfer most of the aerosol problem to peripheral areas.

On considering questions of this sort, it is evident that quantitative data are needed to assess the directions to be taken to control formation

of inorganic aerosol in urban and downwind atmospheres. In addition, the control of anthropogenic inorganic particulates, particularly the stable sulfates, may have worldwide implications with regard to inadvertent climate modification through changes in albedo and by furnishing an excess of water droplet nuclei.

The initial phase of the program to study the formation of sulfate aerosol from SO_2 has been largely completed. The major goals of this portion of the program were to establish the experimental techniques necessary to quantitatively study the loss of SO_2 under simulated atmospheric conditions in the SAPRC all-glass chamber, to characterize the effect of various parameters on SO_2 lifetime and to determine the suitability of using added seed aerosol to facilitate the characterization of aerosol formed from SO_2 .

B. Results and Discussion

Rates of Disappearance of SO_2 . In order to characterize the behavior of SO_2 in the all-glass chamber, a number of control experiments have been carried out. Three types of experiments have been completed: (a) SO_2 disappearance as a function of irradiation intensity and relative humidity, (b) effect of alkene and ozone and of ozone alone on SO_2 lifetime, and (c) loss of SO_2 in the presence of the SAPRC surrogate for ambient pollutants.

The results of experiments to determine the net SO_2 disappearance rate (i.e., corrected for dilution) under the above conditions are given in Table 11. Detailed data sheets for these experiments are presented in Appendix B. Appendix C contains plots of $\ln \text{SO}_2$ versus time showing least squares fits to the experimental data and the total disappearance rates.

As expected⁹¹ the consumption of SO_2 is greatly accelerated in the presence of ozone and propene. The initial (averaged over first half hour) net loss of SO_2 is $\sim 30\% \text{ hr}^{-1}$, but this falls off rapidly to $\sim 0.2\% \text{ hr}^{-1}$ for 1.75 to 5 hours. The decay of SO_2 in the presence of ozone alone was about $\sim 1\% \text{ hr}^{-1}$. The disappearance of SO_2 in the presence of surrogate and NO_x is similar to that observed in pure air at the same humidity during irradiation (see Table 11).

Aerosol Formation. During the latter part of Run 218 at 100% light intensity, propene (and additional SO_2) was added to the reaction mixture

Table 11. Rate of SO₂ Loss in All-Glass Chamber under a Variety of Conditions (using FC-12 as Inert Tracer for Determination of Dilution Rate)

AGC Run No.	RH Range (%)	Amount of Water (torr)	Temp. Range (°C)	Light Intensity	Estimated Net ^a SO ₂ Loss Rate (% hr ⁻¹)
216	1-2.5	0.6-1.0	32.6-33.7	0%	0.4
217 ^b	11-24.5	5.1-10.3	34.4-36.6	100%	3
218 ^c	40-49	15.5-18.9	33.2-33.8	0%	0.4
218 ^d	31-55.5	14.0-22.2	33.0-36.0	100%	5
219 ^e	81.5-71.5	30.1-26.9	32.6-34.2	0%	5
220 ^f	41-40	15.2-15.3	32.7-32.8	0%	30
220 ^g	40-38.5	14.9-14.5	32.8-33.1	0%	0.2
221 ^h	44-38.5	16.4-13.9	32.7-33.2	0%	1 ⁱ
222 ^j	46.3-32.5	15.9-12.7	34.3-30.5	0%	2
222 ^k	32.5-26.0	13.2-12.0	33.8-36.6	100%	4

^aTotal rate of SO₂ disappearance minus dilution rate (as determined from monitoring of FC-12).
^bContinuation of 216 at 100% light intensity; additional SO₂ injected; approx. 0.5 ppm ozone present in this experiment.

^cFlash lamp in Teco 43 replaced; instrument zeroed and spanned using MSA filter and calculated SO₂ in chamber, respectively.

^dOzone increased to ~0.3 ppm in first 6 hours of irradiation and remained at about that value.
^eLamp bases in Teco 43 replaced; zeroed and spanned as in b.

^f1 ppm propene and 0.6 ppm ozone present; average rate for first 30 min of reaction.
^g0.40 ppm propene and 0.20 ppm ozone present; average rate after 1.75 hr of reaction.
^h0.57 ppm ozone added.

ⁱValue for time after ozone was added.

^jSurrogate (~1700 ppbC) and NO (0.3 ppm calc.) present initially.

^kSurrogate (~1100 ppbC) and NO^x (~0.5 ppm) present at beginning of irradiation.

Table 12. Aerosol Measurements Using the Whitby and Climet Analyzers during Propene-Ozone-SO₂ Experiment (AGC-218)

Size (µm)	Concentration (particles cm ⁻³)													
	Initial ^a	1440hrs	1450hrs	1505hrs	1515hrs	1525hrs	1535hrs	1545hrs	1555hrs	1605hrs	1615hrs	1625hrs	1638hrs	1660hrs
<u>Whitby</u>														
.0075	---	---	---	---	---	---	---	---	---	---	---	---	---	---
.0133	---	2.3x10 ³	1.75x10 ⁴	4.17x10 ²	---	---	---	---	---	---	---	---	---	---
.0237	1.17x10 ³	1.72x10 ⁵	1.62x10 ⁵	9.52x10 ⁴	8.85x10 ⁴	6.90x10 ⁴	4.34x10 ⁴	8.35x10 ³	2.71x10 ⁴	1.57x10 ⁴	1.25x10 ⁴	---	---	---
.0422	1.74x10 ²	7.83x10 ²	2.60x10 ⁴	4.09x10 ⁴	2.96x10 ⁴	3.30x10 ⁴	3.65x10 ⁴	4.35x10 ⁴	3.83x10 ⁴	3.44x10 ⁴	2.76x10 ⁴	---	---	---
.0750	1.54x10 ³	1.69x10 ³	5.95x10 ³	1.54x10 ³	2.10x10 ⁴	2.08x10 ⁴	2.26x10 ⁴	2.21x10 ⁴	2.28x10 ⁴	2.46x10 ⁴	2.62x10 ⁴	---	---	---
.133	5.7x10 ³	5.59x10 ³	5.49x10 ³	6.10x10 ³	6.53x10 ³	7.21x10 ³	7.56x10 ³	7.66x10 ³	8.19x10 ³	8.10x10 ³	8.15x10 ³	---	---	---
.237	3.75x10 ³	3.51x10 ³	3.46x10 ³	3.33x10 ³	3.37x10 ³	3.35x10 ³	3.27x10 ³	3.38x10 ³	3.36x10 ³	3.39x10 ³	3.30x10 ³	---	---	---
.422	6.47x10 ²	5.40x10 ²	5.74x10 ²	4.87x10 ²	4.87x10 ²	4.67x10 ²	4.93x10 ²	4.80x10 ²	4.46x10 ²	4.54x10 ²	4.67x10 ²	---	---	---
.750	8.42x10	8.73x10	7.02x10	5.26x10	6.32x10	5.97x10	4.91x10	5.96x10	7.37x10	4.9x10	6.32x10	---	---	---
<u>Climet</u>	Initial ^a	1442hrs	1452hrs	1502hrs	1510hrs	1520hrs	1530hrs	1540hrs	1550hrs	1600hrs	1620hrs	1638hrs		
>.3 ^b	5.0x10	5.7x10	6.28x10	6.78x10	6.78x10	7.24x10	7.52x10	7.83x10	8.35x10	9.15x10	9.11x10	8.95x10		
>.5 ^b	2.3x10 ⁻¹	2.5x10 ⁻¹	3.24x10 ⁻¹	3.24x10 ⁻¹	3.53x10 ⁻¹	1.83x10 ⁻¹	2.97x10 ⁻¹	3.68x10 ⁻¹	3.81x10 ⁻¹	3.11x10 ⁻¹	3.53x10 ⁻¹	3.30x10 ⁻¹		
>1	4.5x10 ⁻²													
>3	1.0x10 ⁻³													

^aPrior to injection at 1424 of SO₂ and propene.

^bDilution system used, 100 to 1.

which contained ~0.3 ppm ozone. Particle size distributions were monitored with both a TSI Electrical Aerosol Analyzer (Whitby) and a Climet Optical Particle Analyzer. Background readings were taken prior to the injection of SO₂. The data obtained with both instruments are given in Table 12. Subsequent work with the Climet has indicated difficulties in calibration and this probably is the source of the variation in results obtained with the two instruments in this study. However, the relative values obtained do show that (1) there was a rapid growth of small particles following the injection of propene in the presence of SO₂ and ozone, and (2) the growth of these particles to larger sizes was accompanied by the decrease of smaller sized particles.

Added Seed Aerosol. A series of calculations were carried out in order to design the most effective experiments for characterizing aerosol stability in the all-glass chamber (AGC). These calculations suggested that the chamber should be seeded with approximately 10^4 cm^{-3} of 0.1 μm diameter particles, a mass loading of about $40 \mu\text{g m}^{-3}$. It was calculated that such a seed cloud should have a life expectancy of approximately 100 hours in the 6400-liter all-glass chamber.

Based on these results, spray-dried K₂SO₄ doped with 10 mole percent of KHSO₄ was used as a test aerosol. This material was intended to play the part of an innocuous surrogate for generated aerosol, organic or inorganic. It was found that the most stable size is in the 0.1 to 0.2 μm diameter range which has an irradiated chamber life expectancy of 30 hours or more (17% or less loss to the walls after 6 hours). Lifetimes estimated from the data decrease rapidly outside of the stable size range, reaching 12 hours at both 0.04 μm and 0.3 μm . An expectancy of 12 hours corresponds to 40% wall loss in 6 hours. These estimates are thought to be conservative estimates representing lower limits on the actual life expectancy.

These findings have the following implications:

- Any experiment involving aerosols, either inorganic or organic, should be seeded if semi-quantitative results on the material yield are desired. The seed particles should be somewhat less than or equal to 0.1 μm in size. Otherwise wall losses can diminish the apparent yield, sometimes to negligibility.
- Measurements of gross light scattering properties like b_{scat} have

no meaning with regard to mass concentration or visibility reduction potential. The particle size distribution resulting from the appearance of aerosol material will be severely deficient in the high light scattering region (0.2 to 1.0 μm). This will be the case for either seeded or unseeded mixtures because of the accumulated wall losses beyond 0.2 μm . In the case of properly seeded mixtures, comparatively few particles will grow into the light scattering region in any case.

Most of the uncertainty about the estimates originates from the limitations of the TSI 3030 (Whitby mobility analyzer). The particle number concentration cannot be reduced to values where the effects of the second order processes of coagulation are negligible during several hours without encountering the sensitivity limits of the TSI 3030. Consequently, the observed decay had to be corrected for coagulation as well as for the exchange rate. For some particle sizes the correction was about equal to the overall decay rate. In addition, methods of estimating coagulation rate constants are subject to considerable uncertainty, especially when the sizes are comparable to the mean free path for air molecules (0.065 μm). The detailed methods of calculation will be presented in our final report.

Consideration of the coagulation rate constants calculated suggests a strategy for obtaining better estimates of wall loss as a function of particle size. The rates imply that coagulation can be dramatically reduced if a monodisperse aerosol is used to test the chamber. Our data suggest that tests with particle sizes around 0.05 μm and 0.45 μm would more definitely determine the loss curve. A test at the larger size would allow use of an optical particle counter to check the TSI 3030 data.

A nearly monodisperse aerosol of size 0.05 μm can be derived from the output of the spray dryer. Figure 8 shows three size distributions of spray-dried aerosols in the AGC according to the TSI 3030. These distributions were measured early in the aerosol's history in order to get as close as possible to the aerosol seen at the spray dryer's output. There is a trend towards larger sizes as the K_2SO_4 concentration is increased. The 7% solution has a good yield near 0.05 μm . Larger sizes can be removed at the cost of considerable loss at 0.05 μm by using a nucleopore filter. Smaller sizes could be removed by either a diffusion battery or by allowing some coagulation in a plastic bag before filtering.




Figure 8. Size Distributions (Whitby) of Spray-Dried Potassium Sulfate Solutions in SAPRC All-Glass Chamber.

V. FURTHER DEVELOPMENT OF AN AIR PARCEL TRANSPORT SIMULATION METHODOLOGY

The purpose of this study was to provide a basis for designing realistic chamber experiments simulating air parcel transport effects during a full day irradiation. In our final report⁹² to ARB Contract No. 4-214, we described in some detail a methodology for determining the primary pollutant concentration profiles in an air parcel as it is transported and gave specific results for an air trajectory which terminated in Pasadena at 1300 Pacific Daylight Time on July 25, 1973. During our current work we have developed procedures for determining the rate of dilution and diffusion for the various pollutants in that air parcel and we describe these here.

As discussed in the previous report⁹² for a given trajectory which is determined by interpolation of ground level meteorological data, the ambient air quality is determined by a similar interpolation procedure using the air monitoring data. Based on the results for the CO concentration history, a diffusion model is generated to allow for the transport of pollutants in the vertical direction. The emissions inventory used in the estimate is that adopted by the General Research Corporation (GRC) from one developed by Systems Applications, Inc.⁹³ and the diffusion model is a portion of the GRC DIFKIN model.⁹⁴ Given these tools, both dilution and fresh pollutant injection rates required in a chamber experiment in order to approximate an air parcel history can then be determined by the following procedure.

The results of the GRC Diffusion program are the concentration profiles in the vertical direction as a function of time of day along the given trajectory of interest. These are based on the diffusion coefficient profiles estimates which yield reasonable CO ground level concentrations as compared with the CO ambient air quality data for that particular day. In order to determine the dilution/diffusion rates for the various pollutants, an air mass has to be defined. In the following example, two air masses were studied--one extended from ground level up to 108 meters, while the other started at 27 meters above ground level and extended up to 135 meters. Note in both cases that the depth of the air parcel was 108 meters. The mass rate of diffusion

transport of the various pollutants was determined from the concentration profile and the diffusion coefficients assumed. The mass flux across the boundary is given by the follow equation:

$$\dot{m} = -D \frac{dc}{dz}$$

where \dot{m} = mass flux rate, D = turbulent diffusion coefficient, c = local concentration, and z = vertical space variable. From this equation the mass flux by diffusion of the pollutants was determined both in and out of the air parcel. In the case where one of the boundaries of the air parcel was the ground level, the source flux was used for the mass flux into the air parcel. In addition to determining the fluxes, the average concentration of pollutants within the parcel was determined by numerically integrating under the concentration profile.

These dynamic processes in the atmosphere must now be related to the chamber operation. We define VDIL as the volume of chamber material which must be removed from the chamber each hour in order to accomplish the efflux of a given pollutant as determined for the air parcel. This amount is computed by the following formula:

$$VDIL = 6400 \left(\frac{\dot{m}_o}{H C_{AVG}} \right)$$

where \dot{m}_o = mass flux out of parcel, H = height of parcel and C_{AVG} = average concentration in parcel. We define VIN as that volume of pure pollutant which then must be added back with the clean air make-up to maintain the chamber conditions. This essentially is the rate of influx predicted from the transport model.

Figures 9 through 12 show the time variation of VDIL and VIN for the two air parcels considered above (i.e., 0-108 m and 27-135 m) for NO_x and for hydrocarbon. It is interesting to note that there is a large variation in these quantities as a function of time of day as one would intuitively expect. The two contributing factors for this large variation are variations in the emissions inventory as a function of time of day




Figure 9. Volume of Chamber Sample to be Removed Each Hour (VDIL) for Simulation of NOx Concentration in Ambient Air Parcels.




Figure 10. Volume of Pure NO_x to be Added to Chamber Each Hour (VIN) for Simulation of NO_x Concentration in Ambient Air Parcels.




Figure 11. Volume of Chamber Sample to be Removed Each Hour (VDIL) for Simulation of Hydrocarbon Concentration in Ambient Air Parcels.




Figure 12. Volume of Pure Hydrocarbon to be Added to Chamber Each Hour (VIN) for Simulation of Hydrocarbon Concentration in Ambient Air Parcels.

and geographic location, and a changing of the diffusion coefficients as the air parcel warms up. Figure 13 shows the variation in the average concentration of the two pollutants as a function of time for the 0-108 m air parcel. It should be noted that for this trajectory there is a continuous increase in the average concentration. Thus, there are no dilution effects apparent. Clearly one would expect that a dilution effect for other trajectories which pass over low source regions later in the day would result.

Recently the choice of dilution rates of approximately 10% per hour by workers operating the large outdoor smog chambers in North Carolina (and by workers in a number of previous chamber studies explicitly employing dilution) has been challenged by Shair^{1,2} as being too low. In essence, our initial data suggest that the situation is complex and that both groups of workers are correct to some extent. For example, based on NO_x , the peak dilution we have calculated is approximately 50%--far above those considered by previous chamber workers. However, for the air parcel beginning at ground level, the average dilution rate over the entire trajectory, again based on NO_x , was 16%; and if one uses only that portion of the trajectory from 6:00 in the morning till 1:00 in the afternoon, the average dilution rate is 21%. Clearly from Figure 11 the dilution rates based on hydrocarbon are considered lower than the above values for NO_x . Thus, the results of this analysis for hydrocarbon appear to be more in line with the lower dilution values used in previous chamber studies.

The fact that dilution rates based on NO_x are considerably larger than those based on hydrocarbon is a crucial observation and results from the fact that the concentration of hydrocarbons is far in excess of the concentration of NO_x . Consequently, less volume need be removed from the chamber to achieve the mass fluxes that occur. Obviously, this situation presents a problem in determining an appropriate simulation in actual chamber experiments and work is continuing to determine what VDIL is required to maintain an appropriate hydrocarbon-to- NO_x ratio, a possibly more realistic control parameter.




Figure 13. Hourly Average Concentrations of NO_x and Hydrocarbons for Ambient Air Parcel.

VI. AMBIENT AIR ANALYSIS

In April, 1976, the SAPRC was selected by the Environmental Protection Agency to conduct a three-year study of trace pollutants in ambient air using a 22-meter base path multiple reflection cell and FT-IR spectrometer which together are capable of operating at total pathlengths in excess of 2 km. This system, designed and initially assembled by P. Hanst and coworkers at the RTP laboratories of EPA, has now been established on the roof of Weber Hall East on the UCR campus several hundred yards from the ARB mobile laboratory and Fawcett Laboratory which houses the SAPRC environmental chamber facility. Since the EPA long-path infrared (LPIR) spectrometer has an effective pathlength which is a factor of 10-20 times that of the in-situ multiple reflection cells in the SAPRC smog chambers, we have chosen to concentrate our efforts in ambient air analysis by LPIR spectroscopy on setting up and operating the 2-km FT-IR spectrometer. This system is now successfully operating and spectra are being recorded routinely (even during the only moderate smog episodes which have been characteristic of this summer). Under SAPRC's policy of resource sharing, results which are obtained using the 2-km FT-IR spectrometer will be considered in the context of the aerosol and gas phase data which are being obtained under our companion ARB contract for investigation of ambient aerosols. Our goal remains to obtain the most complete heterogeneous and homogeneous mass balance for carbon and nitrogen-containing species possible through the combined capabilities of the major instrumental facilities which have now been assembled at SAPRC under the auspices of the ARB, NSF-RANN and EPA,

VII. REFERENCES

1. F. H. Shair, "Photochemical Smog Systems: Heterogeneous Decay of Ozone," Science, 191, 1057 (1976).
2. D. L. Fox, R. Kamens, and H. E. Jeffries, "Photochemical Smog Systems: Heterogeneous Decay of Ozone," Science, 191, 1058 (1976).
3. B. F. Goeller, J. H. Bigelow, J. C. DeHaven, W. T. Mikolowsky, R. L. Petruschell and B. M. Woodfill, "Strategy Alternatives for Oxidant Control in the Los Angeles Region," RAND Corporation R-1368-EPA December, 1973.
4. J. G. French, V. Hasselblad, R. Johnson and K. E. McClain, "The Association of Suspended Nitrates in Ambient Air with the Aggravation of Asthma," preprint provided by J. French (1975).
5. "Sources and Effects of Sulfates in the Atmosphere," California Air Resources Staff Report, March 6, 1975.
6. J. N. Pitts, Jr., P. J. Bekowies, A. M. Winer, G. J. Doyle, J. M. McAfee, and K. W. Wilson, "The Design and Construction of an Environmental Chamber Facility for the Study of Photochemical Air Pollution," Final Report, California ARB Grant No. 5-067-1, in draft.
7. J. N. Pitts, Jr., A. M. Winer, K. R. Darnall, G. J. Doyle, J. M. McAfee, P. J. Bekowies, and W. D. Long, "A Smog Chamber Study of the Role of Hydrocarbons, Oxides of Nitrogen and Aged Smog in the Production of Photochemical Oxidants. I. Facilities, Methodology, and Results for Six-Hour Irradiations," in preparation (1976).
8. "Analysis for Total Oxidant Content of the Atmosphere (2% Potassium Iodide Method)," Recommended Method No. 2A, Air and Industrial Hygiene Laboratory, California State Department of Public Health, Berkeley, California, January, 1969.
9. J. N. Pitts, Jr., J. M. McAfee, W. D. Long, and A. M. Winer, "Long-Path Infrared Spectroscopic Investigation at Ambient Concentrations of the 2% Neutral Buffered Potassium Iodide Method for Determination of Ozone," Environ. Sci. Technol., 10, 787 (1976).
10. State of California Air Resources Board, Sacramento, California, "Comparison of Oxidant Calibration Procedures," Final Report of the Ad Hoc Committee, February 3, 1975.
11. A. M. Winer, J. W. Peters, J. P. Smith, and J. N. Pitts, Jr., "Response of Commercial Chemiluminescent NO-NO₂ Analyzers to Other Nitrogen-Containing Compounds," Environ. Sci. Technol., 8, 1118 (1974).
12. C. W. Spicer and D. F. Miller, "Nitrogen Balance in Smog Chamber Studies," presented at the 67th Annual Meeting of the Air Pollution Control Association, Denver, Colorado, June 1974.

13. E. F. Darley, K. A. Kettner, and E. R. Stephens, "Analysis of Peroxyacyl Nitrates by Gas Chromatography with Electron Capture Detection," Anal. Chem., 35, 589 (1963).
14. E. R. Stephens and M. A. Price, "Analysis of an Important Pollutant: Peroxyacetyl Nitrate," J. Chem. Educ., 50, 351 (1973).
15. R. G. Smith, R. J. Bryan, M. Feldstein, B. Levadie, F. A. Miller, E. R. Stephens, and N. G. White, "Tentative Method of Analysis for Formaldehyde Content of the Atmosphere (Colorimetric Method)," H.L.S., 7, (1) Supplement, 87 (1970).
16. E. R. Stephens and F. R. Burleson, "Distribution of Light Hydrocarbons in Ambient Air," J. Air Pollut. Contr. Assoc., 19, 929 (1969).
17. E. R. Stephens, "Hydrocarbons in Polluted Air," Summary Report, Coordinating Research Council, Project CAPA-5-68, June 1973; NTIS No. PB 230 993/AS.
18. G. J. Doyle, P. J. Bekowies, A. M. Winer, and J. N. Pitts, Jr., "A Charcoal Adsorption Air Purification System for Chamber Studies Investigating Atmospheric Photochemistry," Environ. Sci. Technol., 11, 45 (1977).
19. J. R. Holmes, R. J. O'Brien, J. H. Crabtree, T. A. Hecht, and J. H. Seinfeld, "Measurement of Ultraviolet Radiation Intensity in Photochemical Smog Studies," Environ. Sci. Technol., 7, 519 (1973).
20. G. J. Doyle, A. C. Lloyd, K. R. Darnall, A. M. Winer, and J. N. Pitts, Jr., "Gas Phase Kinetic Study of Relative Rates of Reaction of Selected Aromatic Compounds with Hydroxyl Radicals in an Environmental Chamber," Environ. Sci. Technol., 9, 237 (1975).
21. J. N. Pitts, Jr., A. M. Winer, K. R. Darnall, G. J. Doyle, and J. M. McAfee, "Chemical Consequences of Air Quality Standards and of Control Implementation Programs: Roles of Hydrocarbons, Oxides of Nitrogen, and Aged Smog in the Production of Photochemical Oxidant," Final Report, California Air Resources Board Contract No. 3-017, July, 1975a.
22. A. C. Lloyd, K. R. Darnall, A. M. Winer, and J. N. Pitts, Jr., "Relative Rate Constants for Reaction of the Hydroxyl Radical with a Series of Alkanes, Alkenes, and Aromatic Hydrocarbons," J. Phys. Chem., 80, 789 (1976a).
23. A. C. Lloyd, K. R. Darnall, A. M. Winer, and J. N. Pitts, Jr., "Relative Rate Constants for the Reactions of OH Radicals with Isopropyl Alcohol, Diethyl and Di-n-propyl Ether at 305 ± 2 K," Chem. Phys. Lett., 42, 205 (1976b).
24. A. M. Winer, A. C. Lloyd, K. R. Darnall, and J. N. Pitts, Jr., "Relative Rate Constants for the Reaction of the Hydroxyl Radical with Selected Ketones, Chloroethanes, and Monoterpene Hydrocarbons," J. Phys. Chem., 80, 1635 (1976a).

25. A. M. Winer, A. C. Lloyd, K. R. Darnall, and J. N. Pitts, Jr. "Relative Rate Constants for the Reaction of OH Radicals with Tetrahydrofuran and n-Propyl and sec-Butyl Acetate," Chem. Phys. Lett., submitted (1977).
26. K. R. Darnall, A. M. Winer, A. C. Lloyd, and J. N. Pitts, Jr., "Relative Rate Constants for the Reaction of OH Radicals with Selected C₆ and C₇ Alkanes and Alkenes at 305 ± 2 K," Chem. Phys. Lett., 44, 415 (1976).
27. W. P. L. Carter, A. C. Lloyd, and J. N. Pitts, Jr., unpublished results, 1975.
28. D. H. Stedman and H. Niki, "Ozonolysis Rates of Some Atmospheric Gases," Environ. Lett., 4, 303 (1973).
29. C. T. Pate, R. Atkinson, and J. N. Pitts, Jr., "The Gas Phase Reaction of O₃ with a Series of Aromatic Hydrocarbons," J. Environ. Sci. Health, A11, 3, 1 (1976).
30. R. Atkinson and J. N. Pitts, Jr., "Absolute Rate Constants for the Reaction of O(³P) Atoms with Selected Alkanes, Alkenes, and Aromatics as Determined by a Modulation Technique," J. Phys. Chem., 78, 1780 (1974).
31. D. H. Stedman, C. H. Wu, and H. Niki, "Kinetics of Gas-Phase Reactions of Ozone with some Olefins," J. Phys. Chem., 77, 2511 (1973).
32. S. M. Japar, C. H. Wu, and H. Niki, "Rate Constants for the Reaction of Ozone with Olefins in the Gas Phase," J. Phys. Chem., 78, 2318 (1974).
33. S. M. Japar and H. Niki, "Gas Phase-Reactions of the Nitrate Radical with Olefins," J. Phys. Chem., 79, 1629 (1975).
34. A. C. Lloyd, "Evaluated and Estimated Kinetic Data for Phase Reactions of the Hydroperoxyl Radical," Int. J. Chem. Kinet., 6, 169 (1974).
35. R. A. Perry, R. Atkinson, and J. N. Pitts, Jr., "Rate Constants for the Reaction of OH Radicals with n-butane over the Temperature Range 297-340 K," J. Chem. Phys., 64, 5314 (1976c).
36. N. R. Greiner, "Hydroxyl Radical Kinetics by Kinetic Spectroscopy. VI. Reactions with Alkanes in the Range 300-500 K," J. Chem. Phys., 53, 1070 (1970).
37. I. M. Campbell, B. J. Handy, and R. M. Kirby, "Gas Phase Chain Reactions of H₂O₂ + NO₂ + CO," J. Chem. Soc. Faraday Trans. I, 71, 867 (1975).
38. S. Gordon and W. A. Mulac, "Reaction of the OH (xII) Radical Produced by the Pulse Radiolysis of Water Vapor," Int. J. Chem. Kinet., Symp. No. 1, 289 (1975).

39. R. A. Gorse and D. H. Volman, "Photochemistry of the Gaseous Hydrogen Peroxide-Carbon Monoxide System. II: Rate Constants for Hydroxyl Radical Reactions with Hydrocarbons and for Hydrogen Atom Reactions with Hydrogen Peroxide," J. Photochem., 3, 115 (1974).
40. E. D. Morris, Jr. and H. Niki, "Reactivity of Hydroxyl Radicals with Olefins," J. Phys. Chem., 75, 3640 (1971).
41. F. Stuhl, "Rate Constant for the Reaction of OH with n-C₄H₁₀," Z. Naturforsch., 28A, 1383 (1973).
42. R. Atkinson and J. N. Pitts, Jr., "Rate Constants for the Reaction of OH Radicals with Propylene and the Butenes Over the Temperature Range 297-425°K," J. Chem. Phys., 63, 3591 (1975).
43. N. R. Greiner, "Hydroxyl-Radical Kinetics by Kinetic Spectroscopy. II. Reactions with C₂H₆, C₃H₈, and iso-C₄H₁₀ at 300 K," J. Chem. Phys., 46, 3389 (1967).
44. J. Heicklen, K. Westberg, and N. Cohen, "The Conversion of NO to NO₂ in Polluted Atmospheres," Publication No. 115-69, Center for Air Environmental Studies, University Park, PA, 1969.
45. K. Westberg and N. Cohen, "The Chemical Kinetics of Photochemical Smog as Analyzed by Computer," The Aerospace Corporation, El Segundo, CA ATR-70 (8107)-1. December 30, 1969.
46. B. Weinstock, E. E. Daby, and H. Niki, "Discussion on Paper Presented by E. R. Stephens," in Chemical Reactions in Urban Atmospheres, C. S. Tuesday, Ed., Elsevier, NY, 1971, pp. 54-55.
47. H. Niki, E. E. Daby, and B. Weinstock, "Mechanisms of Smog Reactions," Advan. Chem. Series, 113, 16 (1972).
48. K. L. Demerjian, J. A. Kerr, and J. G. Calvert, "The Mechanism of Photochemical Smog Formation," Advan. Environ. Sci. Technol., 4, 1 (1974).
49. J. N. Pitts, Jr. and B. J. Finlayson, "Mechanisms of Photochemical Air Pollution," Angew. Chem. Int. Ed., 14, I (1975).
50. B. J. Finlayson and J. N. Pitts, Jr., "Photochemistry of the Polluted Troposphere," Science, 192, 111 (1976).
51. C. C. Wang, L. I. Davis, Jr., C. H. Wu, S. Japar, H. Niki, and B. Weinstock, "Hydroxyl Radical Concentrations Measured in Ambient Air," Science, 189, 797 (1975).
52. D. Perner, D. H. Ehhalt, H. W. Patz, U. Platt, E. P. Roth, and A. Volz, "OH Radicals in the Lower Troposphere," 12th International Symposium on Free Radicals, Laguna Beach, CA, January 4-9, 1976.

53. D. D. Davis, W. Heaps, and T. McGee, "Direct Measurements of Natural Tropospheric Levels of OH Via an Aircraft Borne Tunable Dye Laser," Geophys. Res. Lett., 3, 331 (1976).
54. R. Atkinson, D. A. Hansen, and J. N. Pitts, Jr., "Rate Constants for the Reaction of the OH Radical with H₂ and NO(M=Ar and N₂)," J. Chem. Phys., 62, 3284 (1975a).
55. R. Atkinson, D. A. Hansen, and J. N. Pitts, Jr., "Rate Constants for the Reactions of OH radicals with CHF₂Cl, CF₂Cl₂, CFCl₃, and H₂ over the Temperature Range 297-434 K," J. Chem. Phys., 63, 1703 (1975b).
56. D. A. Hansen, R. Atkinson, and J. N. Pitts, Jr., "Rate Constants for the Reaction of OH Radicals with a Series of Aromatic Hydrocarbons," J. Phys. Chem., 79, 1763 (1975).
57. R. A. Perry, R. Atkinson, and J. N. Pitts, Jr., "Rate Constants for the Reaction of OH Radicals with CHFCl₂ and CH₃Cl over the Temperature Range 298-423 K, and with CH₂Cl₂ at 298 K," J. Chem. Phys., 64, 1618 (1976a).
58. R. A. Perry, R. Atkinson, and J. N. Pitts, Jr., "Rate Constants for the Reactions OH + H₂S → H₂O + SH and OH + NH₃ → H₂O + NH₂ Over the Temperature Range 297-427 K," J. Chem. Phys., 64, 3237 (1976b).
59. R. Atkinson, R. A. Perry, and J. N. Pitts, Jr., "Rate Constants for the Reaction of OH Radicals with 2-methyl-2-butene over the temperature range 297-425 K," Chem. Phys. Lett., 38, 607 (1976a).
60. R. Atkinson, R. A. Perry, and J. N. Pitts, Jr., "Rate Constants for the Reactions of the OH Radical with NO₂ (M=Ar and N₂) and SO₂ (M=Ar)," J. Chem. Phys., 65, 306 (1976b).
61. R. Atkinson, R. A. Perry, and J. N. Pitts, Jr., "Rate Constants for the Reaction of OH Radicals with Ethylene Over the Temperature Range 299-425°K," J. Chem. Phys., Submitted for publication (1976c).
62. R. Atkinson, K. R. Darnall, A. C. Lloyd, A. M. Winer, and J. N. Pitts, Jr., "Kinetics and Mechanisms of the Reactions of the Hydroxyl Radical with Organic Compounds in the Gas Phase," Accts. Chem. Res., submitted for publication (1976).
63. C. H. Wu, S. M. Japar, and H. Niki, "Relative Reactivities of HO-Hydrocarbon Reactions from Smog Reactor Studies," J. Environ. Sci. Health, A11, 191 (1976).
64. W. A. Glasson and C. S. Tuesday, "Hydrocarbon Reactivity in the Atmospheric Photooxidation of Nitric Oxide," paper presented at the 150th National Meeting, ACS, Atlantic City, NJ, September, 1965.
65. R. F. Hampson, Jr. and D. Garvin, Eds., "Chemical Kinetic and Photochemical Data for Modeling Atmospheric Chemistry," NBS Technical Note 866, June, 1975.

66. K. R. Darnall, A. C. Lloyd, A. M. Winer, and J. N. Pitts, Jr., "Reactivity Scale for Atmospheric Hydrocarbons Based on Reaction with Hydroxyl Radical," Environ. Sci. Technol., 10, 692 (1976a).
67. J. N. Pitts, Jr., A. C. Lloyd, A. M. Winer, K. R. Darnall, and G. J. Doyle, "Development and Application of a Hydrocarbon Reactivity Scale Based on Reaction with the Hydroxyl Radical," Paper No. 76-31.1, 69th Annual Air Pollution Control Association Meeting, Portland, Oregon, June 27-July 1, 1976c.
68. J. N. Pitts, Jr., "Reactivity Scale for Atmospheric Hydrocarbons Based on Reaction with Hydroxyl Radical," presentation to California Air Resources Board Committee on Photochemical Reactivity, December, 1975.
69. J. N. Pitts, Jr., A. M. Winer, K. R. Darnall, A. C. Lloyd, and G. J. Doyle, "Hydrocarbon Reactivity and the Role of Hydrocarbons, Oxides of Nitrogen and Aged Smog in the Production of Photochemical Oxidants," International Conference on Photochemical Oxidant Pollution and Its Control, Raleigh, NC, September 13-17, 1976d.
70. A. P. Altshuller and J. J. Bufalini, "Photochemical Aspects of Air Pollution: A Review," Environ. Sci. Technol., 5, 39 (1971).
71. A. P. Altshuller and I. R. Cohen, "Structural Effects on the Rate of Nitrogen Dioxide Formation in the Photooxidation of Organic Compound-Nitric Oxide Mixtures in Air," Int. J. Air Water Pollut., 7, 787 (1963).
72. E. A. Schuck and G. J. Doyle, "Photooxidation of Hydrocarbons in Mixtures Containing Oxides of Nitrogen and Sulfur Dioxide," Report No. 29, Air Pollution Foundation, San Marino, CA, 1959.
73. E. R. Stephens and W. E. Scott, "Relative Reactivity of Various Hydrocarbons in Polluted Atmospheres," Proc. Amer. Petrol. Inst., 42 (III), 665 (1962).
74. C. S. Tuesday, "Atmospheric Photooxidation of Olefins. Effect of Nitrogen Oxides," Arch. Environ. Health, 7, 188 (1963).
75. A. P. Altschuller, I. R. Cohen, S. F. Sleva, and S. L. Kopczynski "Air Pollution: Photooxidation of Aromatic Hydrocarbons," Science, 135, 442 (1962).
76. S. L. Kopczynski, "Photooxidation of Alkylbenzene-Nitrogen Dioxide Mixtures in Air," Int. J. Air Water Pollut., 8, 107 (1964).
77. S. L. Kopczynski, in A. P. Altshuller and J. J. Bufalini, "Photochemical Aspects of Air Pollution: A Review," Environ. Sci. Technol., 5, 39 (1971).
78. EPA-650/3-74-010, "Proceedings of the Solvent Reactivity Conference," U. S. Environmental Protection Agency, Research Triangle Park, NC, November, 1974.

79. ARB, "Adoption of a System for the Classification of Organic Compounds According to Photochemical Reactivity," California Air Resources Board Staff Report No. 76-3-4, February 19, 1976a.
80. ARB, "Board Approves Reactivity Classification," California Air Resources Board Bulletin, 7, No. 3, March, 1976b, p. 4.
81. H. Niki, P. Maker, C. Savage, and L. Breitenbach, "Fourier Transform Spectroscopic Studies of Organic Species Participating in Photochemical Smog Formation," International Conference on Environmental Sensing and Assessment, Las Vegas, Nevada, September 14-19, 1975.
82. W. P. L. Carter, K. R. Darnall, A. C. Lloyd, A. M. Winer, and J. N. Pitts, Jr., "Evidence for Alkoxy Radical Isomerization in Photooxidations of C₄-C₆ Alkanes Under Simulated Atmospheric Conditions," Chem. Phys. Lett., 42, 22 (1976b).
83. K. R. Darnall, W. P. L. Carter, A. M. Winer, A. C. Lloyd, and J. N. Pitts, Jr., "Importance of RO₂ + NO in Alkyl Nitrate Formation from C₄-C₆ Alkane Photooxidations Under Simulated Atmospheric Conditions," J. Phys. Chem., 80, 1948 (1976c).
84. R. J. Charlson, A. H. Vanderpol, D. S. Covert, A. P. Waggoner, and N. C. Ahlquist, "Sulfuric Acid - Ammonium Sulfate Aerosol: Optical Detection in the St. Louis Region," Science, 184, (4133), 156 (1974).
85. J. Barone, T. A. Cahill, R. G. Flocchini and D. J. Shadoan, "Visibility Reduction: A Characterization of Three Urban Sites in California," submitted to Science (1975).
86. D. Grosjean and S. K. Friedlander, "Gas-Particle Distribution Factors for Organic and Other Pollutants in the Los Angeles Atmosphere," submitted for Publication, presented in part at 67th Annual Air Pollution Control Association Meeting, Denver, Colorado, June 9-13, 1974.
87. "Conference on Health Effects of Atmospheric Salts and Gases of Sulfur and Nitrogen in Association with Photochemical Oxidant," Newport Beach, California, January 7-8, 1974, Vols. I and II, T. T. Crocker, Editor.
88. R. S. Chapman, C. M. Shy, J. F. Finklea, D. E. House, H. E. Goldberg and C. G. Hayes, "Chronic Respiratory Disease," Arch. Env. Health, 27 (3), 138-142 (1973).
89. J. G. French, G. Lowimore, W. C. Nelson, J. F. Finklea, T. English, and M. H. Hertz, "The Effect of Sulfur Dioxide and Suspended Sulfates on Acute Respiratory Disease," Arch. Env. Health, 27 (3), 129 (1973).
90. G. M. Hidy, "Characterization of Aerosols in California (ACHEX)," Final Report, Volume 1, Summary, submitted to California Air Resources Board, September 30, 1974.

91. R. A. Cox and S. A. Penkett, "Aerosol Formation from Sulphur Dioxide in the Presence of Ozone and Olefinic Hydrocarbons," J. Chem. Soc. Faraday Trans. I, 68, 1735 (1972).
92. J. N. Pitts, Jr., A. M. Winer, K. R. Darnall, G. J. Doyle, and J. M. McAfee, "Chemical Consequences of Air Quality Standards and of Control Implementation Programs: Roles of Hydrocarbons, Oxides of Nitrogen, and Aged Smog in the Production of Photochemical Oxidant," Final Report, California Air Resources Board Contract No. 4-214, May, 1976a.
93. P. J. W. Roberts, P. M. Roth, and C. L. Nelson, "Contaminant Emissions in the Los Angeles Basin--Their Sources, Rates, and Distribution," Appendix A of Development of a Simulation Model for Estimating Ground Level Concentrations of Photochemical Pollutants, Systems Applications, Inc., Report No. 71SAI-6, March, 1971.
94. A. Q. Eschenroeder and J. R. Martinez, "Concepts and Applications of Photochemical Smog Models," in Photochemical Smog and Ozone Reactions, Advan. Chem. Series, 113, 101 (1972).

VIII. LIST OF PUBLICATIONS FROM SAPRC-ARB CHAMBER PROGRAM

"A Novel 20 KW Solar Simulator Designed for Air Pollution Research," J. H. Beauchene, P. J. Bekowies, J. M. McAfee, A. M. Winer, L. Zafonte, and J. N. Pitts, Jr., Proceedings of Seventh Conference on Space Simulation (NASA) Special Publication 336, Los Angeles, California, November 12-14, 1973, Paper No. 66, pp. 811-825.

"Response of Commercial Chemiluminescent NO-NO₂ Analyzers to other Nitrogen Containing Compounds," A. M. Winer, J. W. Peters, J. P. Smith, and J. N. Pitts, Jr., Environ. Sci. Technol., 8, 1118 (1974).

"Experimentally Validated Mechanisms, Models and Control Strategies for Photochemical Air Pollution," J. N. Pitts, Jr., W. P. Carter, K. R. Darnall, G. J. Doyle, W. Kuby, A. C. Lloyd, J. M. McAfee, C. Pate, J. P. Smith, J. L. Sprung, and A. M. Winer, Proceedings of the Second Annual NSF-RANN Trace Contaminants Conference, Asilomar, California, August 29-31, 1974, LBL-3217, pp. 50-53.

"A Gas Phase Kinetic Study of the Relative Rates of Reaction of Selected Aromatic Compounds, with Hydroxyl Radicals in an Environmental Chamber," G. J. Doyle, A. C. Lloyd, K. R. Darnall, A. M. Winer, and J. N. Pitts, Jr., Environ. Sci. Technol., 9, 237 (1975).

"Long-Path Infrared Spectroscopic Investigation at Ambient Concentrations of the 2% Neutral Buffered Potassium Iodide Method for Determination of Ozone," J. N. Pitts, Jr., J. M. McAfee, W. D. Long, and A. M. Winer, Environ. Sci. Technol., 10, 787 (1976).

"Infrared Absorptivity of the 9.6 μ m Ozone Band as a Function of Spectral Resolution and Abundance," J. M. McAfee, E. R. Stephens, D. R. Fitz, and J. N. Pitts, Jr., J. Quant. Spectrosc. Radiat. Transfer, 16, 829 (1976).

"Relative Rate Constants for Reaction of the Hydroxyl Radical with a Series of Alkanes, Alkenes, and Aromatic Hydrocarbons," A. C. Lloyd, K. R. Darnall, A. M. Winer, and J. N. Pitts, Jr., J. Phys. Chem., 80, 789 (1976).

"Relative Rate Constants for the Reaction of the Hydroxyl Radical with Selected Ketones, Chloroethenes, and Monoterpene Hydrocarbons," A. M. Winer, A. C. Lloyd, K. R. Darnall, and J. N. Pitts, Jr., J. Phys. Chem., 80, 1635 (1976).

"Reactivity Scale for Atmospheric Hydrocarbons Based on Reaction with Hydroxyl Radical," K. R. Darnall, A. C. Lloyd, A. M. Winer, and J. N. Pitts, Jr., Environ. Sci. Technol., 10, 692 (1976).

"Development and Application of a Hydrocarbon Reactivity Scale Based on Reaction with the Hydroxyl Radical," J. N. Pitts, Jr., A. C. Lloyd, A. M. Winer, K. R. Darnall, and G. J. Doyle, 69th Annual Air Pollution Control Association Meeting, Portland, Oregon, June 27-July 1, 1976, Paper No. 76-31.1.

"Relative Rate Constants for the Reactions of OH Radicals with Isopropyl Alcohol, Diethyl and Di-n-Propyl Ether at 305 ± 2 K," A. C. Lloyd, K. R. Darnall, A. M. Winer, and J. N. Pitts, Jr., Chem. Phys. Lett., 42, 205 (1976).

"Hydrocarbon Reactivity and the Role of Hydrocarbons, Oxides of Nitrogen, and Aged Smog in the Production of Photochemical Oxidants, J. N. Pitts, Jr., A. M. Winer, K. R. Darnall, A. C. Lloyd, and G. J. Doyle, Proceedings of the International Conference on Photochemical Oxidant Pollution and Its Control, Raleigh, North Carolina, Sept. 13-17, 1976, EPA-600/3-77-0016, January 1977, pp. 687-704.

"Relative Rate Constants for the Reaction of OH Radicals with Selected C₆ and C₇ Alkanes and Alkenes at 305 ± 2 K," K. R. Darnall, A. M. Winer, A. C. Lloyd, and J. N. Pitts, Jr., Chem. Phys. Lett., 44, 415 (1976).

"Charcoal-Adsorption Air Purification System for Chamber Studies Investigating Atmospheric Photochemistry," G. J. Doyle, P. J. Bekowies, A. M. Winer, and J. N. Pitts, Jr., Environ. Sci. Technol., 11, 45 (1977).

"Atmospheric Chemical and Physical Transformations of Oxide of Nitrogen," J. N. Pitts, Jr., D. Grosjean, A. M. Winer, K. R. Darnall, and G. J. Doyle, State of California Air Resources Board Photochemical/Transport Workshop, University of California, Los Angeles, California, January 6-7, 1977.